

2-LOCAL TRIPLE DERIVATIONS ON VON NEUMANN ALGEBRAS

KARIMBERGEN KUDAYBERGENOV, TIMUR OIKHBERG,
ANTONIO M. PERALTA, AND BERNARD RUSSO

ABSTRACT. We prove that every (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra M is a triple derivation, equivalently, the set $\text{Der}_t(M)$, of all triple derivations on M , is algebraically 2-reflexive in the set $\mathcal{M}(M) = M^M$ of all mappings from M into M .

1. INTRODUCTION

Let X and Y be Banach spaces. According to the terminology employed in the literature (see, for example, [4]), a subset \mathcal{D} of the Banach space $B(X, Y)$, of all bounded linear operators from X into Y , is called *algebraically reflexive* in $B(X, Y)$ when it satisfies the property:

$$(1.1) \quad T \in B(X, Y) \text{ with } T(x) \in \mathcal{D}(x), \forall x \in X \Rightarrow T \in \mathcal{D}.$$

Algebraic reflexivity of \mathcal{D} in the space $L(X, Y)$, of all linear mappings from X into Y , a stronger version of the above property not requiring continuity of T , is defined by:

$$(1.2) \quad T \in L(X, Y) \text{ with } T(x) \in \mathcal{D}(x), \forall x \in X \Rightarrow T \in \mathcal{D}.$$

In 1990, Kadison proved that (1.1) holds if \mathcal{D} is the set $\text{Der}(M, X)$ of all (associative) derivations on a von Neumann algebra M into a dual M -bimodule X [18]. Johnson extended Kadison's result by establishing that the set $\mathcal{D} = \text{Der}(A, X)$, of all (associative) derivations from a C^* -algebra A into a Banach A -bimodule X satisfies (1.2) [17].

Algebraic reflexivity of the set of local triple derivations on a C^* -algebra and on a JB^* -triple have been studied in [24, 9, 12] and [14]. More precisely, Mackey proves in [24] that the set $\mathcal{D} = \text{Der}_t(M)$, of all triple derivations on a JBW^* -triple M satisfies (1.1). The result has been supplemented in [12], where Burgos, Fernández-Polo and the third author of this note prove

2010 *Mathematics Subject Classification*. Primary 46L05; 46L40.

Key words and phrases. triple derivation; 2-local triple derivation.

Second author partially supported by Simons Foundation travel grant 210060. Third author partially supported by the Spanish Ministry of Science and Innovation, D.G.I. project no. MTM2011-23843, Junta de Andalucía grant FQM375, and the Deanship of Scientific Research at King Saud University (Saudi Arabia) research group no. RG-1435-020.

that for each JB*-triple E , the set $\mathcal{D} = \text{Der}_t(E)$ of all triple derivations on E satisfies (1.2).

Hereafter, *algebraic reflexivity* will refer to the stronger version (1.2) which does not assume the continuity of T .

In [6], Brešar and Šemrl proved that the set of all (algebra) automorphisms of $B(H)$ is algebraically reflexive whenever H is a separable, infinite-dimensional Hilbert space. Given a Banach space X . A linear mapping $T : X \rightarrow X$ satisfying the hypothesis at (1.2) for $\mathcal{D} = \text{Aut}(X)$, the set of automorphisms on X , is called a *local automorphism*. Larson and Sourour showed in [22] that for every infinite dimensional Banach space X , every surjective local automorphism T on the Banach algebra $B(X)$, of all bounded linear operators on X , is an automorphism.

Motivated by the results of Šemrl in [31], references witness a growing interest in a subtle version of algebraic reflexivity called *algebraic 2-reflexivity* (cf. [1, 2, 10, 11, 21, 23, 25, 26] and [29]). A subset \mathcal{D} of the set $\mathcal{M}(X, Y) = Y^X$, of all mappings from X into Y , is called *algebraically 2-reflexive* when the following property holds: for each mapping T in $\mathcal{M}(X, Y)$ such that for each $a, b \in X$, there exists $S = S_{a,b} \in \mathcal{D}$ (depending on a and b), with $T(a) = S_{a,b}(a)$ and $T(b) = S_{a,b}(b)$, then T lies in \mathcal{D} . A mapping $T : X \rightarrow Y$ satisfying that for each $a, b \in X$, there exists $S = S_{a,b} \in \mathcal{D}$ (depending on a and b), with $T(a) = S_{a,b}(a)$ and $T(b) = S_{a,b}(b)$ will be called a *2-local \mathcal{D} -mapping*. If we assume that every mapping $S \in \mathcal{D}$ is r -homogeneous (that is, $S(ta) = t^r S(a)$ for every $t \in \mathbb{R}$ or \mathbb{C}) with $0 < r$, then every 2-local \mathcal{D} -mapping $T : X \rightarrow Y$ is r -homogeneous. Indeed, for each $a \in X$, $t \in \mathbb{C}$ take $S_{a,ta} \in \mathcal{D}$ satisfying $T(ta) = S_{a,ta}(ta) = t^r S_{a,ta}(a) = t^r T(a)$.

Šemrl establishes in [31] that for every infinite-dimensional separable Hilbert space H , the sets $\text{Aut}(B(H))$ and $\text{Der}(B(H))$, of all (algebra) automorphisms and associative derivations on $B(H)$, respectively, are algebraically 2-reflexive in $\mathcal{M}(B(H)) = \mathcal{M}(B(H), B(H))$. Ayupov and the first author of this note proved in [1] that the same statement remains true for general Hilbert spaces (see [20] for the finite dimensional case). Actually, the set $\text{Hom}(A)$, of all homomorphisms on a general C*-algebra A , is algebraically 2-reflexive in the Banach algebra $B(A)$, of all bounded linear operators on A , and the set ${}^*\text{-Hom}(A)$, of all * -homomorphisms on A , is algebraically 2-reflexive in the space $L(A)$, of all linear operators on A (cf. [27]).

In recent contributions, Burgos, Fernández-Polo and the third author of this note prove that the set ${}^*\text{-Hom}(M)$ (respectively, $\text{Hom}_t(M)$), of all * -homomorphisms (respectively, triple homomorphisms) on a von Neumann algebra (respectively, on a JBW*-triple) M , is an algebraically 2-reflexive subset of $\mathcal{M}(M)$ (cf. [10], [11], respectively), while Ayupov and the first author of this note establish that set $\text{Der}(M)$ of all derivations on M is algebraically 2-reflexive in $\mathcal{M}(M)$ (see [2]).

In this paper, we consider the set $\text{Der}_t(A)$ of all triple derivations on a C^* -algebra A . We recall that every C^* -algebra A can be equipped with a ternary product of the form

$$\{a, b, c\} = \frac{1}{2}(ab^*c + cb^*a).$$

When A is equipped with this product it becomes a JB^* -triple in the sense of [19]. A linear mapping $\delta : A \rightarrow A$ is said to be a *triple derivation* when it satisfies the (triple) Leibnitz rule:

$$\delta\{a, b, c\} = \{\delta(a), b, c\} + \{a, \delta(b), c\} + \{a, b, \delta(c)\}.$$

It is known that every triple derivation is automatically continuous (cf. [3]). We refer to [3, 15] and [28] for the basic references on triple derivations. According to the standard notation, 2-local $\text{Der}_t(A)$ -mappings from A into A are called *2-local triple derivations*.

The goal of this note is to explore the algebraic 2-reflexivity of $\text{Der}_t(A)$ in $\mathcal{M}(A)$. Our main result proves that every (not necessarily linear nor continuous) 2-local triple derivation on an arbitrary von Neumann algebra M is a triple derivation (hence linear and continuous) (see Theorem 2.14), equivalently, $\text{Der}_t(M)$ is algebraically 2-reflexive in $\mathcal{M}(M)$.

2. 2-LOCAL TRIPLE DERIVATIONS ON VON NEUMANN ALGEBRAS

We start by recalling some generalities on triple derivations. Let A be a C^* -algebra. For each $b \in A$, we shall denote by M_b the Jordan multiplication mapping by the element b , that is $M_b(x) = b \circ x = \frac{1}{2}(bx + xb)$. Following standard notation, given elements a, b in A , we denote by $L(a, b)$ the operator on A defined by $L(a, b)(x) = \{a, b, x\} = \frac{1}{2}(ab^*x + xb^*a)$. It is known that the mapping $\delta(a, b) : A \rightarrow A$, given by

$$\delta(a, b)(x) = L(a, b)(x) - L(b, a)(x),$$

is a triple derivation on A (cf. [3, 15]). A triple derivation which is a finite linear combination of derivations of the form $\delta(a, b)$ is called an *inner triple derivation*.

Let $\delta : A \rightarrow A$ be a triple derivation on a unital C^* -algebra. By [15, Lemmas 1 and 2], $\delta(\mathbf{1})^* = -\delta(\mathbf{1})$, and $M_{\delta(\mathbf{1})} = \delta(\frac{1}{2}\delta(\mathbf{1}), \mathbf{1})$ is an inner triple derivation on A and the difference $D = \delta - \delta(\frac{1}{2}\delta(\mathbf{1}), \mathbf{1})$ is a Jordan $*$ -derivation on A , more concretely,

$$D(x \circ y) = D(x) \circ y + x \circ D(y), \text{ and } D(x^*) = D(x)^*,$$

for every $x, y \in A$. By [3, Corollary 2.2], δ (and hence D) is a continuous operator. A widely known result, due to B.E. Johnson, states that every bounded Jordan derivation from a C^* -algebra A to a Banach A -bimodule is an associative derivation (cf. [16]). Therefore, D is an associative $*$ -derivation in the usual sense. When $A = M$ is a von Neumann algebra, we can guarantee that D is an inner derivation, that is there exists $\tilde{a} \in A$ satisfying $D(x) = [\tilde{a}, x] = \tilde{a}x - x\tilde{a}$, for every $x \in A$ (cf. [30, Theorem

4.1.6]). Further, from the condition $D(x^*) = D(x)^*$, for every $x \in A$, we deduce that $(\tilde{a}^* + \tilde{a})x = x(\tilde{a}^* + \tilde{a})$. Thus, taking $a = \frac{1}{2}(\tilde{a} - \tilde{a}^*)$, it follows that $[a, x] = [\tilde{a}, x]$, for every $x \in M$. We have therefore shown that for every triple derivation δ on a von Neumann algebra M , there exist skew-hermitian elements $a, b \in M$ satisfying

$$\delta(x) = [a, x] + b \circ x,$$

for every $x \in M$.

Our first lemma is a direct consequence of the above arguments (see [15, Lemmas 1 and 2]).

Lemma 2.1. *Let $T : A \rightarrow A$ be a (not necessarily linear nor continuous) 2-local triple derivation on a unital C^* -algebra. Then*

- (a) $T(\mathbf{1})^* = -T(\mathbf{1})$;
- (b) $M_{T(\mathbf{1})} = \delta(\frac{1}{2}T(\mathbf{1}), \mathbf{1})$ is an inner triple derivation on A ;
- (c) $\hat{T} = T - \delta(\frac{1}{2}T(\mathbf{1}), \mathbf{1})$ is a 2-local triple derivation on A with $\hat{T}(\mathbf{1}) = 0$.

□

In what follows, we denote by A_{sa} the hermitian elements of the C^* -algebra A .

Lemma 2.2. *Let $T : A \rightarrow A$ be a (not necessarily linear nor continuous) 2-local triple derivation on a unital C^* -algebra satisfying $T(\mathbf{1}) = 0$. Then $T(x) = T(x)^*$ for all $x \in A_{sa}$.*

Proof. Let $x \in A_{sa}$. By assumptions,

$$\begin{aligned} T(x)^* &= \{\mathbf{1}, T(x), \mathbf{1}\} = \{\mathbf{1}, \delta_{x, \mathbf{1}}(x), \mathbf{1}\} = \delta_{x, \mathbf{1}}\{\mathbf{1}, x, \mathbf{1}\} - 2\{\delta_{x, \mathbf{1}}(\mathbf{1}), x, \mathbf{1}\} \\ &= \delta_{x, \mathbf{1}}(x^*) - 2\{T(\mathbf{1}), x, \mathbf{1}\} = \delta_{x, \mathbf{1}}(x) = T(x). \end{aligned}$$

The proof is complete. □

Lemma 2.3. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra satisfying $T(\mathbf{1}) = 0$. Then for every $x, y \in M_{sa}$ there exists a skew-hermitian element $a_{x,y} \in M$ such that*

$$T(x) = [a_{x,y}, x], \text{ and, } T(y) = [a_{x,y}, y].$$

Proof. For every $x, y \in M_{sa}$ we can find skew-hermitian elements $a_{x,y}, b_{x,y} \in M$ such that

$$T(x) = [a_{x,y}, x] + b_{x,y} \circ x, \text{ and, } T(y) = [a_{x,y}, y] + b_{x,y} \circ y.$$

Taking into account that $T(x) = T(x)^*$ (see Lemma 2.2) we obtain

$$\begin{aligned} [a_{x,y}, x] + b_{x,y} \circ x &= T(x) = T(x)^* = [a_{x,y}, x]^* + (b_{x,y} \circ x)^* \\ &= [x, a_{x,y}^*] + x \circ b_{x,y}^* = [x, -a_{x,y}] - x \circ b_{x,y} = [a_{x,y}, x] - b_{x,y} \circ x, \end{aligned}$$

i.e. $b_{x,y} \circ x = 0$, and similarly $b_{x,y} \circ y = 0$. Therefore $T(x) = [a_{x,y}, x]$, $T(y) = [a_{x,y}, y]$, and the proof is complete. □

We state now an observation, which plays an useful role in our study.

Lemma 2.4. *Let a and b be skew-hermitian elements in a C^* -algebra A . Suppose $x \in A$ is self-adjoint with $[a, x] + 2b \circ x = 0$. Then $[a, x] = 0$ and $b \circ x = 0$.*

Proof. Since $0 = ax - xa + bx + xb$. Passing to the adjoint, we obtain $ax - xa - (bx + xb) = 0$. Conclude the proof by adding and subtracting these two equalities. The proof is complete. \square

Let M be a von Neumann algebra. If $x \in M_{sa}$, we denote by $s(x)$ the support projection of x – that is, the projection onto $(\ker(x))^\perp = \overline{\text{ran}(x)}$. We say that x has full support if $s(x) = 1$ (equivalently, $\ker(x) = \{0\}$).

Lemma 2.5. *Let M be a von Neumann algebra. Suppose $u \in M_+$ has full support, $c \in M$ is self-adjoint, and $\sigma(c^2u) \cap (0, \infty) = \emptyset$. Then $c = 0$. Consequently, if u and c are as above, and $uc + cu = 0$ (or $c^2u = -cuc \leq 0$), then $c = 0$.*

Proof. For the first statement of the lemma, suppose $\sigma(c^2u) \cap (0, \infty) = \emptyset$. Note that

$$(-\infty, 0] \supseteq \sigma(c^2u) \cup \{0\} = \sigma(c \cdot cu) \supseteq \sigma(cuc).$$

However, cuc is positive, hence $\sigma(cuc) \subset [0, \|cuc\|]$, with $\max_{\lambda \in \sigma(cuc)} = \|cuc\|$. Thus, $cu^{1/2}u^{1/2}c = cuc = 0$, which means that $cu^{1/2} = u^{1/2}c = 0$ and hence $s(c) \leq 1 - s(u^{1/2}) = 1 - s(u) = 0$, which leads to $c = 0$.

To prove the second part, we have $c^2u = -cuc \leq 0$, hence in particular, $\sigma(c^2u) \subset (-\infty, 0]$. The proof is complete. \square

In [2, Lemma 2.2], Ayupov and the first author of this note prove that for every (not necessarily linear nor continuous) 2-local derivation on a von Neumann algebra $\Delta : M \rightarrow M$, and every self-adjoint element $z \in M$, there exists $a \in M$ satisfying

$$\Delta(x) = [a, x],$$

for every $x \in \mathcal{W}^*(z)$, where $\mathcal{W}^*(z) = \{z\}''$ denotes the abelian von Neumann subalgebra of M generated by the element z , and the unit element and $\{z\}''$ denotes the bicommutant of the set $\{z\}$. We prove next a ternary version of this result.

Lemma 2.6. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra. Let $z \in M$ be a self-adjoint element and let $\mathcal{W}^*(z) = \{z\}''$ be the abelian von Neumann subalgebra of M generated by the element z and the unit element. Then there exist skew-hermitian elements $a_z, b_z \in M$, depending on z , such that*

$$T(x) = [a_z, x] + b_z \circ x = a_z x - x a_z + \frac{1}{2}(b_z x + x b_z)$$

for all $x \in \mathcal{W}^*(z)$. In particular, T is linear on $\mathcal{W}^*(z)$.

Proof. We can assume that $z \neq 0$. Note that the abelian von Neumann subalgebras generated by $\mathbf{1}$ and z and by $\mathbf{1}$ and $\mathbf{1} + \frac{z}{2\|z\|}$ coincide. So, replacing z with $\mathbf{1} + \frac{z}{2\|z\|}$ we can assume that z is an invertible positive element.

By definition, there exist skew-hermitian elements $a_z, b_z \in M$ (depending on z) such that

$$T(z) = [a_z, z] + b_z \circ z.$$

Define a mapping $T_0 : M \rightarrow M$ given by $T_0(x) = T(x) - ([a_z, z] + b_z \circ z)$, $x \in M$. Clearly, T_0 is a 2-local triple derivation on M . We shall show that $T_0 \equiv 0$ on $\mathcal{W}^*(z)$. Let $x \in \mathcal{W}^*(z)$ be an arbitrary element. By assumptions, there exist skew-hermitian elements $c_{z,x}, d_{z,x} \in M$ such that

$$T_0(z) = [c_{z,x}, z] + d_{z,x} \circ z, \text{ and, } T_0(x) = [c_{z,x}, x] + d_{z,x} \circ x.$$

Since $0 = T_0(z) = [c_{z,x}, z] + d_{z,x} \circ z$, we get $[c_{z,x}, z] + d_{z,x} \circ z = 0$.

Taking into account that z is a hermitian element and Lemma 2.4 we get $c_{z,x}z = zc_{z,x}$ and $d_{z,x}z = -zd_{z,x}$.

Since z has a full support, and $d_{z,x}^2z = -d_{z,x}zd_{z,x}$, Lemma 2.5 implies that $d_{z,x} = 0$. Further

$$c_{z,x} \in \{z\}' = \{z\}''' = \mathcal{W}^*(z)',$$

i.e. $c_{z,x}$ commutes with any element in $\mathcal{W}^*(z)$. Therefore $T_0(x) = [c_{z,x}, x] + d_{z,x} \circ x = 0$, for all $x \in \mathcal{W}^*(z)$. The proof is complete. \square

2.1. Complete additivity of 2-local derivations and 2-local triple derivations on von Neumann algebras.

Let $\mathcal{P}(M)$ denote the lattice of projections in a von Neumann algebra M . Let X be a Banach space. A mapping $\mu : \mathcal{P}(M) \rightarrow X$ is said to be *finitely additive* when

$$\mu \left(\sum_{i=1}^n p_i \right) = \sum_{i=1}^n \mu(p_i),$$

for every family p_1, \dots, p_n of mutually orthogonal projections in M . A mapping $\mu : \mathcal{P}(M) \rightarrow X$ is said to be *bounded* when the set

$$\{\|\mu(p)\| : p \in \mathcal{P}(M)\}$$

is bounded.

The celebrated Bunce-Wright-Mackey-Gleason theorem ([7, 8]) states that if M has no summand of type I_2 , then every bounded finitely additive mapping $\mu : \mathcal{P}(M) \rightarrow X$ extends to a bounded linear operator from M to X .

According to the terminology employed in [32] and [13], a completely additive mapping $\mu : \mathcal{P}(M) \rightarrow \mathbb{C}$ is called a *charge*. The Dorofeev–Sherstnev theorem ([32, Theorem 29.5] or [13, Theorem 2]) states that any charge on a von Neumann algebra with no summands of type I_n is bounded.

We shall use the Dorofeev-Shertsnev theorem in Corollary 2.8 in order to be able to apply the Bunce-Wright-Mackey-Gleason theorem in Proposition 2.9. To this end, we need Proposition 2.7, which is implicitly applied in [2, proof of Lemma 2.3] for 2-local associative derivations. A proof is included here for completeness reasons.

First, we recall some facts about the strong* topology. For each normal positive functional φ in the predual of a von Neumann algebra M , the mapping

$$x \mapsto \|x\|_\varphi = \left(\varphi\left(\frac{xx^* + x^*x}{2}\right) \right)^{\frac{1}{2}} \quad (x \in M)$$

defines a prehilbertian seminorm on M . The *strong* topology* of M is the locally convex topology on M defined by all the seminorms $\|\cdot\|_\varphi$, where φ runs in the set of all positive functionals in M_* (cf. [30, Definition 1.8.7]). It is known that the strong* topology of M is compatible with the duality (M, M_*) , that is a functional $\psi : M \rightarrow \mathbb{C}$ is strong* continuous if and only if it is weak* continuous (see [30, Corollary 1.8.10]). We also recall that the product of every von Neumann algebra is jointly strong* continuous on bounded sets (see [30, Proposition 1.8.12]).

Suppose $X = W$ is another von Neumann algebra, and let τ denote the norm, the weak* or the strong* topology of W . The mapping μ is said to be τ -completely additive (respectively, countably or sequentially τ -additive) when

$$(2.1) \quad \mu \left(\sum_{i \in I} p_i \right) = \tau\text{-} \sum_{i \in I} \mu(p_i)$$

for every family (respectively, sequence) $\{p_i\}_{i \in I}$ of mutually orthogonal projections in M .

It is known that every family $(p_i)_{i \in I}$ of mutually orthogonal projections in a von Neumann algebra M is summable with respect to the weak* topology of M and $p = \text{weak}^*\text{-} \sum_{i \in I} p_i$ is a projection in M (cf. [30, Definition 1.13.4]).

Further, for each normal positive functional ϕ in M_* and every finite set $F \subset I$, we have

$$\left\| p - \sum_{i \in F} p_i \right\|_\phi^2 = \phi \left(p - \sum_{i \in F} p_i \right),$$

which implies that the family $(p_i)_{i \in I}$ is summable with respect to the strong* topology of M with the same limit, that is, $p = \text{strong}^*\text{-} \sum_{i \in I} p_i$.

Proposition 2.7. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra. Then the following statements hold:*

- (a) *The restriction $T|_{\mathcal{P}(M)}$ is sequentially strong* additive, and consequently sequentially weak* additive;*
 (b) *$T|_{\mathcal{P}(M)}$ is weak* completely additive, i.e.,*

$$(2.2) \quad T \left(\text{weak}^* \text{-} \sum_{i \in I} p_i \right) = \text{weak}^* \text{-} \sum_{i \in I} T(p_i)$$

for every family $(p_i)_{i \in I}$ of mutually orthogonal projections in M .

Proof. (a) Let $(p_n)_{n \in \mathbb{N}}$ be a sequence of mutually orthogonal projections in M . Let us consider the element $z = \sum_{n \in \mathbb{N}} \frac{1}{n} p_n$. By Lemma 2.6 there exist skew-hermitian elements $a_z, b_z \in M$ such that $T(x) = [a_z, x] + b_z \circ x$ for all $x \in \mathcal{W}^*(z)$. Since $\sum_{n=1}^{\infty} p_n, p_m \in \mathcal{W}^*(z)$, for all $m \in \mathbb{N}$, and the product of M is jointly strong* continuous, we obtain that

$$\begin{aligned} T \left(\sum_{n=1}^{\infty} p_n \right) &= \left[a_z, \sum_{n=1}^{\infty} p_n \right] + b_z \circ \left(\sum_{n=1}^{\infty} p_n \right) \\ &= \sum_{n=1}^{\infty} [a_z, p_n] + \sum_{n=1}^{\infty} b_z \circ p_n = \sum_{n=1}^{\infty} T(p_n), \end{aligned}$$

i.e. $T|_{\mathcal{P}(M)}$ is a countably or sequentially strong* additive mapping.

(b) Let φ be a positive normal functional in M_* , and let $\|\cdot\|_{\varphi}$ denote the prehilbertian seminorm given by $\|z\|_{\varphi}^2 = \frac{1}{2} \varphi(zz^* + z^*z)$ ($z \in M$). Let $\{p_i\}_{i \in I}$ be an arbitrary family of mutually orthogonal projections in M . For every $n \in \mathbb{N}$ define

$$I_n = \{i \in I : \|T(p_i)\|_{\varphi} \geq 1/n\}.$$

We claim, that I_n is a finite set for every natural n . Otherwise, passing to a subset if necessary, we can assume that there exists a natural k such that I_k is infinite and countable. In this case the series $\sum_{i \in I_k} T(p_i)$ does not converge with respect to the semi-norm $\|\cdot\|_{\varphi}$. On the other hand, since I_k is a countable set, by (a), we have

$$T \left(\sum_{i \in I_k} p_i \right) = \text{strong}^* \text{-} \sum_{i \in I_k} T(p_i),$$

which is impossible. This proves the claim.

We have shown that the set

$$I_0 = \left\{ i \in I : \|T(p_i)\|_{\varphi} \neq 0 \right\} = \bigcup_{n \in \mathbb{N}} I_n$$

is a countable set, and $\|T(p_i)\|_{\varphi} = 0$, for every $i \in I \setminus I_0$.

Set $p = \sum_{i \in I \setminus I_0} p_i \in M$. We shall show that $\varphi(T(p)) = 0$. Let q denote the support projection of φ in M . Having in mind that $\|T(p_i)\|_\varphi^2 = 0$, for every $i \in I \setminus I_0$, we deduce that $T(p_i) \perp q$ for every $i \in I \setminus I_0$.

Replacing T with $\widehat{T} = T - \delta(\frac{1}{2}T(\mathbf{1}), \mathbf{1})$ we can assume that $T(\mathbf{1}) = 0$ (cf. Lemma 2.1) and $T(x) = T(x)^*$, for every $x \in M_{sa}$ (cf. Lemma 2.2). By Lemma 2.3, for every $i \in I \setminus I_0$ there exists a skew-hermitian element $a_i = a_{p,p_i} \in M$ such that

$$T(p) = a_i p - p a_i, \text{ and, } T(p_i) = a_i p_i - p_i a_i.$$

Since $T(p_i) \perp q$ we get $(a_i p_i - p_i a_i)q = q(a_i p_i - p_i a_i) = 0$, for all $i \in I \setminus I_0$. Thus, since $p a_i p_i q = p_i a_i q$,

$$\begin{aligned} (T(p)p_i)q &= (a_i p - p a_i)p_i q = a_i p_i q - p a_i p_i q \\ &= a_i p_i q - p_i a_i q = (a_i p_i - p_i a_i)q = 0, \end{aligned}$$

and similarly

$$q(p_i T(p)) = 0,$$

for every $i \in I \setminus I_0$. Consequently,

$$(2.3) \quad (T(p)p)q = T(p) \left(\sum_{i \in I \setminus I_0} p_i \right) q = 0 = q \left(\sum_{i \in I \setminus I_0} p_i \right) T(p) = q(pT(p)).$$

Therefore,

$$\begin{aligned} T(p) &= \delta_{p,\mathbf{1}}(p) = \delta_{p,\mathbf{1}}\{p, p, p\} = 2\{\delta_{p,\mathbf{1}}(p), p, p\} + \{p, \delta_{p,\mathbf{1}}(p), p\} \\ &= 2\{T(p), p, p\} + \{p, T(p), p\} = pT(p) + T(p)p + pT(p)^*p \\ &= pT(p) + T(p)p + pT(p)p, \end{aligned}$$

which implies that

$$\begin{aligned} \varphi(T(p)) &= \varphi(pT(p) + T(p)p + pT(p)p) \\ &= \varphi(qpT(p)q) + \varphi(qT(p)pq) + \varphi(qpT(p)pq) = (\text{by (2.3)}) = 0. \end{aligned}$$

Finally, by (a) we have

$$T \left(\sum_{i \in I_0} p_i \right) = \|\cdot\|_{\varphi^-} \sum_{i \in I_0} T(p_i).$$

Two more applications of (a) give:

$$\begin{aligned} \varphi \left(T \left(\sum_{i \in I} p_i \right) \right) &= \varphi \left(T \left(p + \sum_{i \in I_0} p_i \right) \right) = \varphi \left(T(p) + T \left(\sum_{i \in I_0} p_i \right) \right) \\ &= \varphi(T(p)) + \varphi \left(T \left(\sum_{i \in I_0} p_i \right) \right) = \sum_{i \in I_0} \varphi(T(p_i)). \end{aligned}$$

By the Cauchy-Schwarz inequality, $0 \leq |\varphi T(p_i)|^2 \leq \|T(p_i)\|_\varphi^2 = 0$, for every $i \in I \setminus I_0$, and hence $\sum_{i \in I_0} \varphi(T(p_i)) = \sum_{i \in I} \varphi(T(p_i))$. The arbitrariness of φ shows that $T\left(\text{weak}^*\text{-}\sum_{i \in I} p_i\right) = \text{weak}^*\text{-}\sum_{i \in I} T(p_i)$. \square

Let ϕ be a normal functional in the predual of a von Neumann algebra M . Our previous Proposition 2.7 assures that for every (not necessarily linear nor continuous) 2-local triple derivation $T : M \rightarrow M$ the mapping $\phi \circ T|_{\mathcal{P}(M)} : \mathcal{P}(M) \rightarrow \mathbb{C}$ is a completely additive mapping or a charge on M . Under the additional hypothesis of M being a continuous von Neumann algebra or, more generally, a von Neumann algebra with no Type I_n -factors ($1 < n < \infty$) direct summands (i.e. without direct summand isomorphic to a matrix algebra $M_n(\mathbb{C})$, $1 < n < \infty$), the Dorofeev–Sherstnev theorem ([32, Theorem 29.5] or [13, Theorem 2]) imply that $\phi \circ T|_{\mathcal{P}(M)}$ is a bounded charge, that is, the set $\{|\phi \circ T(p)| : p \in \mathcal{P}(M)\}$ is bounded. The uniform boundedness principle gives:

Corollary 2.8. *Let M be a von Neumann algebra with no Type I_n -factor direct summands ($1 < n < \infty$) and let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation. Then the restriction $T|_{\mathcal{P}(M)}$ is a bounded weak* completely additive mapping.* \square

2.2. Additivity of 2-local triple derivations on hermitian parts of von Neumann algebras.

Suppose now that M is a von Neumann algebra with no Type I_n -factor direct summands ($1 < n < \infty$), and $T : M \rightarrow M$ is a (not necessarily linear nor continuous) 2-local triple derivation. By Corollary 2.8 combined with the Bunce-Wright-Mackey-Gleason theorem [7, 8], there exists a bounded linear operator $G : M \rightarrow M$ satisfying that $G(p) = T(p)$, for every projection $p \in M$.

Let z be a self-adjoint element in M . By Lemma 2.6, there exist skew-hermitian elements $a_z, b_z \in M$ such that $T(x) = [a_z, x] + b_z \circ x$, for every $x \in \mathcal{W}^*(z)$. Since $G|_{\mathcal{W}^*(z)}, T|_{\mathcal{W}^*(z)} : \mathcal{W}^*(z) \rightarrow M$ are bounded linear operators, which coincide on the set of projections of $\mathcal{W}^*(z)$, and every self-adjoint element in $\mathcal{W}^*(z)$ can be approximated in norm by finite linear combinations of mutually orthogonal projections in $\mathcal{W}^*(z)$, it follows that $T(x) = G(x)$ for every $x \in \mathcal{W}^*(z)$, and hence

$$T(a) = G(a), \text{ for every } a \in M_{sa},$$

in particular, T is additive on M_{sa} .

The above arguments materialize in the following result.

Proposition 2.9. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra with no Type*

I_n -factor direct summands ($1 < n < \infty$). Then the restriction $T|_{M_{sa}}$ is additive. \square

Corollary 2.10. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a properly infinite von Neumann algebra. Then the restriction $T|_{M_{sa}}$ is additive.*

Next we shall show that the conclusion of the above corollary is also true for a finite von Neumann algebra.

First we show that every 2-local triple derivation on a von Neumann algebra “intertwines” central projections.

Lemma 2.11. *If T is a (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra M , and p is a central projection in M , then $T(Mp) \subset Mp$. In particular, $T(px) = pT(x)$ for every $x \in M$.*

Proof. Consider $x \in Mp$, then $x = pxp = \{x, p, p\}$. T coincides with a triple derivation $\delta_{x,p}$ on the set $\{x, p\}$, hence

$$T(x) = \delta_{x,p}(x) = \delta_{x,p}\{x, p, p\} = \{\delta_{x,p}(x), p, p\} + \{x, \delta_{x,p}(p), p\} + \{x, p, \delta_{x,p}(p)\}$$

lies in Mp .

For the final statement, fix $x \in M$, and consider skew-hermitian elements $a_{x,xp}, b_{x,xp} \in M$ satisfying

$$T(x) = [a_{x,xp}, x] + b_{x,xp} \circ x, \text{ and } T(xp) = [a_{x,xp}, xp] + b_{x,xp} \circ (xp).$$

The assumption p being central implies that $pT(x) = T(px)$. \square

Proposition 2.12. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a finite von Neumann algebra. Then the restriction $T|_{M_{sa}}$ is additive.*

Proof. Since M is finite there exists a faithful normal semi-finite trace τ on M . We shall consider the following two cases.

Case 1. Suppose τ is a finite trace. Replacing T with $\widehat{T} = T - \delta(\frac{1}{2}T(\mathbf{1}), \mathbf{1})$ we can assume that $T(\mathbf{1}) = 0$ (cf. Lemma 2.1) and $T(x) = T(x)^*$, for every $x \in M_{sa}$ (cf. Lemma 2.2). By Lemma 2.3, for every $x, y \in M_{sa}$ there exists a skew-hermitian element $a_{x,y} \in M$ such that $T(x) = [a_{x,y}, x]$ and $T(y) = [a_{x,y}, y]$. Then

$$T(x)y + xT(y) = [a_{x,y}, x]y + x[a_{x,y}, y] = [a_{x,y}, xy],$$

that is,

$$[a_{x,y}, xy] = T(x)y + xT(y).$$

Further

$$0 = \tau([a_{x,y}, xy]) = \tau(T(x)y + xT(y)),$$

i.e. $\tau(T(x)y) = -\tau(xT(y))$, for every $x, y \in M_{sa}$. For arbitrary $u, v, w \in M_{sa}$, set $x = u + v$, and $y = w$. The above identity implies

$$\tau(T(u+v)w) = -\tau((u+v)T(w)) =$$

$= -\tau(uT(w)) - \tau(vT(w)) = \tau(T(u)w) + \tau(T(v)w) = \tau((T(u) + T(v))w)$,
and so

$$\tau((T(u+v) - T(u) - T(v))w) = 0$$

for all $u, v, w \in M_{sa}$. Take $w = T(u+v) - T(u) - T(v)$. Then $\tau(ww^*) = 0$. Since the trace τ is faithful it follows that $ww^* = 0$, and hence $w = 0$. Therefore $T(u+v) = T(u) + T(v)$.

Case 2. As in *Case 1*, we may assume $T(\mathbf{1}) = 0$. Suppose now that τ is a semi-finite trace. Since M is finite there exists a family of mutually orthogonal central projections $\{z_i\}$ in M such that z_i has finite trace for all i and $\bigvee z_i = \mathbf{1}$ (cf. [30, §2.2 or Corollary 2.4.7]). By Lemma 2.11, for each i , T maps z_iM into itself. From Case 1, $T|_{z_iM} : z_iM \rightarrow z_iM$ is additive. Furthermore,

$$z_iT(x+y) = T|_{z_iM}(z_ix + z_iy) = T|_{z_iM}(z_ix) + T|_{z_iM}(z_iy) = z_iT(x) + z_iT(y),$$

for every $x, y \in M$ and every i . Therefore

$$\begin{aligned} T(x+y) &= \left(\sum_i z_i \right) T(x+y) = \sum_i z_i T(x+y) = \sum_i (z_i T(x) + z_i T(y)) \\ &= \left(\sum_i z_i \right) T(x) + \left(\sum_i z_i \right) T(y) = T(x) + T(y), \end{aligned}$$

for every $x, y \in M$. The proof is complete. \square

Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on an arbitrary von Neumann algebra. In this case there exist orthogonal central projections $z_1, z_2 \in M$ with $z_1 + z_2 = \mathbf{1}$ such that:

- (–) z_1M is a finite von Neumann algebra;
 - (–) z_2M is a properly infinite von Neumann algebra,
- (cf. [30, §2.2]).

By Lemma 2.11, for each $k = 1, 2$, z_kT maps z_kM into itself. By Corollary 2.10 and Proposition 2.12 both z_1T and z_2T are additive on M_{sa} . So $T = z_1T + z_2T$ also is additive on M_{sa} .

We have thus proved the following result:

Proposition 2.13. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on an arbitrary von Neumann algebra. Then the restriction $T|_{M_{sa}}$ is additive.* \square

2.3. Main result.

We can state now the main result of this section.

Theorem 2.14. *Let M be an arbitrary von Neumann algebra and let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation. Then T is a triple derivation (hence linear and continuous). Equivalently, the set $Der_t(M)$, of all triple derivations on M , is algebraically 2-reflexive in the set $\mathcal{M}(M) = M^M$ of all mappings from M into M .*

We need the following two Lemmata.

Lemma 2.15. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra with $T(\mathbf{1}) = 0$. Then there exists a skew-hermitian element $a \in M$ such that $T(x) = [a, x]$, for all $x \in M_{sa}$.*

Proof. Let $x \in M_{sa}$. By Lemma 2.3 there exist a skew-hermitian element $a_{x,x^2} \in M$ such that $T(x) = [a_{x,x^2}, x]$, $T(x^2) = [a_{x,x^2}, x^2]$.

Thus,

$$T(x^2) = [a_{x,x^2}, x^2] = [a_{x,x^2}, x]x + x[a_{x,x^2}, x] = T(x)x + xT(x),$$

i.e.

$$(2.4) \quad T(x^2) = T(x)x + xT(x),$$

for every $x \in M_{sa}$.

By Proposition 2.13 and Lemma 2.2, $T|_{M_{sa}} : M_{sa} \rightarrow M_{sa}$ is a real linear mapping. Now, we consider the linear extension \hat{T} of $T|_{M_{sa}}$ to M defined by

$$\hat{T}(x_1 + ix_2) = T(x_1) + iT(x_2), \quad x_1, x_2 \in M_{sa}.$$

Taking into account the homogeneity of T , Proposition 2.13 and the identity (2.4) we obtain that \hat{T} is a Jordan derivation on M . By [5, Theorem 1] any Jordan derivation on a semi-prime algebra is a derivation. Since M is von Neumann algebra, \hat{T} is a derivation on M (see also [33] and [16]). Therefore there exists an element $a \in M$ such that $\hat{T}(x) = [a, x]$ for all $x \in M$. In particular, $T(x) = [a, x]$ for all $x \in M_{sa}$. Since $T(M_{sa}) \subseteq M_{sa}$, we can assume that $a^* = -a$, which completes the proof. \square

Lemma 2.16. *Let $T : M \rightarrow M$ be a (not necessarily linear nor continuous) 2-local triple derivation on a von Neumann algebra. If $T|_{M_{sa}} \equiv 0$, then $T \equiv 0$.*

Proof. Let $x \in M$ be an arbitrary element and let $x = x_1 + ix_2$, where $x_1, x_2 \in M_{sa}$. Since T is homogeneous, by passing to the element $(1 + \|x_2\|)^{-1}x$ if necessary, we can suppose that $\|x_2\| < 1$. In this case the element $y = \mathbf{1} + x_2$ is positive and invertible. Take skew-hermitian elements $a_{x,y}, b_{x,y} \in M$ such that

$$T(x) = [a_{x,y}, x] + b_{x,y} \circ x, \quad \text{and} \quad T(y) = [a_{x,y}, y] + b_{x,y} \circ y.$$

Since $T(y) = 0$, we get $[a_{x,y}, y] + b_{x,y} \circ y = 0$. By Lemma 2.4 we obtain that $[a_{x,y}, y] = 0$ and $ib_{x,y} \circ y = 0$. Taking into account that $ib_{x,y}$ is hermitian, y is positive and invertible, Lemma 2.5 implies that $b_{x,y} = 0$.

We further note that $0 = [a_{x,y}, y] = [a_{x,y}, \mathbf{1} + x_2] = [a_{x,y}, x_2]$, i.e. $[a_{x,y}, x_2] = 0$. Now, $T(x) = [a_{x,y}, x] + b_{x,y} \circ x = [a_{x,y}, x_1 + ix_2] = [a_{x,y}, x_1]$, i.e. $T(x) = [a_{x,y}, x_1]$. Therefore,

$$T(x)^* = [a_{x,y}, x_1]^* = [x_1, a_{x,y}^*] = [x_1, -a_{x,y}] = [a_{x,y}, x_1] = T(x).$$

So $T(x)^* = T(x)$. Now, replacing x by ix we obtain, from the homogeneity of T , that $T(x)^* = -T(x)$. Combining the last two identities we obtain that $T(x) = 0$, which finishes the proof. \square

Proof of Theorem 2.14. Let us define $\widehat{T} = T - \delta\left(\frac{1}{2}T(\mathbf{1}), \mathbf{1}\right)$. Then \widehat{T} is a 2-local triple derivation on M with $\widehat{T}(\mathbf{1}) = 0$ (cf. Lemma 2.1) and $\widehat{T}(x) = \widehat{T}(x)^*$, for every $x \in M_{sa}$ (cf. Lemma 2.2). By Lemma 2.15 there exists an element $a \in M$ such that $\widehat{T}(x) = [a, x]$ for all $x \in M_{sa}$. Consider the 2-local triple derivation $\widehat{T} - [a, \cdot]$. Since $(\widehat{T} - [a, \cdot])|_{M_{sa}} \equiv 0$, Lemma 2.16 implies that $\widehat{T} = [a, \cdot]$, and hence $T = [a, \cdot] + \delta\left(\frac{1}{2}T(\mathbf{1}), \mathbf{1}\right)$, witnessing the desired statement. \square

REFERENCES

- [1] Sh. Ayupov, K.K. Kudaybergenov, 2-local derivations and automorphisms on $B(H)$, *J. Math. Anal. Appl.* **395**, no. 1, 15-18 (2012).
- [2] Sh. Ayupov, K.K. Kudaybergenov, 2-local derivations on von Neumann algebras, to appear in *Positivity*. DOI 10.1007/s11117-014-0307-3.
- [3] T.J. Barton, Y. Friedman, Bounded derivations of JB*-triples, *Quart. J. Math. Oxford* **41**, 255-268 (1990).
- [4] C. Batty, L. Molnar, On topological reflexivity of the groups of *-automorphisms and surjective isometries of $B(H)$, *Arch. Math.* **67**, 415-421 (1996).
- [5] M. Brešar, Jordan derivations on semiprime rings, *Proc. Amer. Math. Soc.* **104**, 1003-1006 (1988).
- [6] M. Brešar, P. Šemrl, On local automorphisms and mappings that preserve idempotents, *Studia Math.* **113**, no. 2, 101-108 (1995).
- [7] L.J. Bunce, J.D.M. Wright, The Mackey-Gleason problem, *Bull. Amer. Math. Soc.* **26**, 288-293 (1992).
- [8] L.J. Bunce, J.D.M. Wright, The Mackey-Gleason problem for vector measures on projections in von Neumann algebras, *J. London Math. Soc.* **49**, 133-149 (1994).
- [9] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, A.M. Peralta, Local triple derivations on C*-algebras, *Communications in Algebra* **42**, 1276-1286 (2014).
- [10] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, A.M. Peralta, A Kowalski-Słodkowski theorem for 2-local *-homomorphisms on von Neumann algebras, preprint 2014.
- [11] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, A.M. Peralta, 2-local triple homomorphisms on von Neumann algebras and JBW*-triples, preprint 2014.
- [12] M. Burgos, F.J. Fernández-Polo, A.M. Peralta, Local triple derivations on C*-algebras and JB*-triples, *Bull. London Math. Soc.*, doi:10.1112/blms/bdu024. arXiv:1303.4569v2.
- [13] S. Dorofeev, On the problem of boundedness of a signed measure on projections of a von Neumann algebra, *J. Funct. Anal.* **103**, 209-216 (1992).
- [14] F.J. Fernández-Polo, A. Molino Salas, A.M. Peralta, Local triple derivations on real C*-algebras and JB*-triples, to appear in *Bull. Malaysian Math. Sci. Soc.*
- [15] T. Ho, J. Martinez-Moreno, A.M. Peralta, B. Russo, Derivations on real and complex JB*-triples, *J. London Math. Soc. (2)* **65**, no. 1, 85-102 (2002).
- [16] B.E. Johnson, Symmetric amenability and the nonexistence of Lie and Jordan derivations, *Math. Proc. Cambridge Philos. Soc.* **120**, no. 3, 455-473 (1996).
- [17] B.E. Johnson, Local derivations on C*-algebras are derivations, *Trans. Amer. Math. Soc.* **353**, 313-325 (2001).
- [18] R.V. Kadison, Local derivations, *J. Algebra* **130**, 494-509 (1990).

- [19] W. Kaup, A Riemann mapping theorem for bounded symmetric domains in complex Banach spaces, *Math. Z.* **183**, 503-529 (1983).
- [20] S.O. Kim, J.S. Kim, Local automorphisms and derivations on \mathbb{M}_n , *Proc. Amer. Math. Soc.* **132**, no. 5, 1389-1392 (2004).
- [21] S.O. Kim, J.S. Kim, Local automorphisms and derivations on certain C^* -algebras, *Proc. Amer. Math. Soc.* **133**, no. 11, 3303-3307 (2005).
- [22] D.R. Larson and A.R. Sourour, Local derivations and local automorphisms of $B(X)$, *Proc. Sympos. Pure Math.* **51**, Part 2, Providence, Rhode Island 1990, pp. 187-194.
- [23] J.-H. Liu, N.-C. Wong, 2-Local automorphisms of operator algebras, *J. Math. Anal. Appl.* **321** 741-750 (2006).
- [24] M. Mackey, Local derivations on Jordan triples, *Bull. London Math. Soc.* **45**, no. 4, 811-824 (2013). doi: 10.1112/blms/bdt007
- [25] L. Molnar, 2-local isometries of some operator algebras, *Proc. Edinburgh Math. Soc.* **45**, 349-352 (2002).
- [26] L. Molnar, Local automorphisms of operator algebras on Banach spaces, *Proc. Amer. Math. Soc.* **131**, 1867-1874 (2003).
- [27] A.M. Peralta, A note on 2-local representations of C^* -algebras, to appear in *Operators and Matrices*.
- [28] A. M. Peralta and B. Russo, Automatic continuity of triple derivations on C^* -algebras and JB^* -triples, *Journal of Algebra* **399**, 960-977 (2014).
- [29] F. Pop, On local representation of von Neumann algebras, *Proc. Amer. Math. Soc.* **132**, No. 12, Pages 3569-3576 (2004).
- [30] S. Sakai, *C^* -algebras and W^* -algebras*, Springer-Verlag, Berlin 1971.
- [31] P. Šemrl, Local automorphisms and derivations on $B(H)$, *Proc. Amer. Math. Soc.* **125**, 2677-2680 (1997).
- [32] A.N. Sherstnev, *Methods of bilinear forms in noncommutative theory of measure and integral*, Moscow, Fizmatlit, 2008, 256 pp.
- [33] A.M. Sinclair, Jordan homomorphisms and derivations on semisimple Banach algebras, *Proc. Amer. Math. Soc.* (3) **24**, 209-214 (1970).

E-mail address: karim2006@mail.ru

CH. ABDIROV 1, DEPARTMENT OF MATHEMATICS, KARAKALPAK STATE UNIVERSITY,
NUKUS 230113, UZBEKISTAN

E-mail address: oikhberg@illinois.edu

DEPT. OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA IL 61801, USA

E-mail address: oikhberg@illinois.edu

E-mail address: aperalta@ugr.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD
DE GRANADA, 18071 GRANADA, SPAIN.

Current address: Visiting Professor at Department of Mathematics, College of Science,
King Saud University, P.O.Box 2455-5, Riyadh-11451, Kingdom of Saudi Arabia.

E-mail address: brusso@uci.edu

DEPARTMENT OF MATHEMATICS, UC IRVINE, IRVINE CA, USA