

ALGEBRAS WHOSE RIGHT NUCLEUS IS A CENTRAL SIMPLE ALGEBRA

S. PUMPLÜN

ABSTRACT. We generalize Amitsur's construction of central simple algebras which are split by transcendental extensions to nonassociative algebras: for every central division algebra D over a field F of characteristic zero there exists an infinite-dimensional unital nonassociative algebra whose right nucleus is D and whose left and middle nucleus are a transcendental extension of F splitting D .

We then give a short direct proof that every p -algebra of degree m , which has a purely inseparable splitting field K of degree m and exponent one, is a differential extension of K and cyclic.

We obtain finite-dimensional division algebras over a field F of characteristic $p > 0$ whose right nucleus is a division p -algebra.

INTRODUCTION

In 1954, Amitsur [2] observed that all associative central division algebras over a field F of characteristic zero can be constructed using differential polynomials. His construction method can be considered as an analogue to the well known crossed product construction, except that he uses the transcendental splitting fields of the algebras instead of their algebraic splitting fields. Some of his results also work for p -algebras, i.e. over base fields of characteristic $p > 0$.

In this paper, we consider algebras which are also obtained from differential polynomials, but which are nonassociative.

These algebras are constructed using the differential polynomial ring $K[t; \delta]$, where K is a field and δ a derivation on K and were defined by Petit [12]: Given a differential polynomial $f \in K[t; \delta]$ of degree m , the set of all differential polynomials of degree less than m , together with the addition given by the usual addition of polynomials, can be equipped with a nonassociative ring structure using right division by f to define the multiplication $g \circ h = gh \bmod_r f$. The resulting nonassociative unital ring S_f , also denoted by $K[t; \delta]/K[t; \delta]f$, is an algebra over the field of constants $F = \text{Const}(\delta)$ of δ . If f generates a two-sided ideal in $K[t; \delta]$, then S_f is the (associative) quotient algebra obtained by factoring out the two-sided principal ideal generated by f .

Date: 25.6.2016.

1991 Mathematics Subject Classification. Primary: 17A35; Secondary: 17A60, 16S36.

Key words and phrases. Differential polynomial ring, skew polynomial, differential polynomial, differential operator, differential algebra, nonassociative algebra, right nucleus.

If f is not two-sided and δ not trivial, then the nuclei of S_f are larger than the center $F = \text{Const}(\delta)$. Then the left and middle nucleus are always given by K , whereas the right nucleus reflects both the choice of f and the structure of the ring $K[t; \delta]$.

We proceed as follows: The basic terminology and notation we use can be found in [2] and Section 1. Section 2 collects some results for those algebras S_f which have a central simple algebra as their right nucleus. For this we employ Amitsur's A -polynomials. In Sections 3 and 4 we show how to construct algebras S_f with a given central simple algebra as right nucleus, first for base fields of characteristic zero, then for base fields of characteristic p : for every central simple algebra B of degree m over a field F of characteristic zero which is split by a transcendental field extension K/F in which F is algebraically closed, there exists an infinite-dimensional unital algebra $S_f = K[t; \delta]/K[t; \delta]f$ over F with right nucleus B (and left and middle nucleus K), see Theorem 8. In particular, there exists an infinite-dimensional unital algebra S_f with right nucleus D for every central division algebra D over F (Corollary 9).

We give a short proof that every p -algebra B of degree m over a field F of characteristic p which is split by a purely inseparable field extension K/F of exponent and degree m is isomorphic to a differential extension (K, δ, d_0) of K (Theorem 13), only invoking a result on the structure of S_f and Amitsur's result [2, Lemma 20]. Thus it is cyclic by [8, Main Theorem].

For every division p -algebra D of degree m over a field F of characteristic p which is split by a purely inseparable field extension K/F of exponent one such that $m < [K : F]$, there is a unital division algebra $S_f = K[t; \delta]/K[t; \delta]f$ over F of dimension mp^e with right nucleus D and left and middle nucleus K (Theorem 15 and Corollary 18). The smallest possible dimension l of such a division algebra containing D as right nucleus is bounded via $m^2 < l \leq mp^{m-1}$ and connected to the number of cyclic algebras that are needed when expressing D as a product of cyclic algebras of degree p in the Brauer group $Br(F)$ (Corollary 18).

1. PRELIMINARIES

1.1. Nonassociative algebras. Let F be a field and let A be an F -vector space. A is an *algebra* over F if there exists an F -bilinear map $A \times A \rightarrow A$, $(x, y) \mapsto x \cdot y$, denoted simply by juxtaposition xy , the *multiplication* of A . An algebra A is called *unital* if there is an element in A , denoted by 1, such that $1x = x1 = x$ for all $x \in A$. We will only consider unital algebras from now on without explicitly saying so.

An algebra $A \neq 0$ is called a *division algebra* if for any $a \in A$, $a \neq 0$, the left multiplication with a , $L_a(x) = ax$, and the right multiplication with a , $R_a(x) = xa$, are bijective. If A has finite dimension over F , A is a division algebra if and only if A has no zero divisors [15, pp. 15, 16].

Associativity in A is measured by the *associator* $[x, y, z] = (xy)z - x(yz)$. The *left nucleus* of A is defined as $\text{Nuc}_l(A) = \{x \in A \mid [x, A, A] = 0\}$, the *middle nucleus* of A is $\text{Nuc}_m(A) = \{x \in A \mid [A, x, A] = 0\}$ and the *right nucleus* of A as $\text{Nuc}_r(A) = \{x \in A \mid [A, A, x] = 0\}$. $\text{Nuc}_l(A)$, $\text{Nuc}_m(A)$, and $\text{Nuc}_r(A)$ are associative subalgebras of A . Their

intersection $\text{Nuc}(A) = \{x \in A \mid [x, A, A] = [A, x, A] = [A, A, x] = 0\}$ is the *nucleus* of A . $\text{Nuc}(A)$ is an associative subalgebra of A containing $F1$ and $x(yz) = (xy)z$ whenever one of the elements x, y, z is in $\text{Nuc}(A)$. The *center* of A is $C(A) = \{x \in A \mid x \in \text{Nuc}(A) \text{ and } xy = yx \text{ for all } y \in A\}$.

1.2. Differential polynomial rings. Let K be a field and $\delta : K \rightarrow K$ a *derivation*, i.e. an additive map such that

$$\delta(ab) = a\delta(b) + \delta(a)b$$

for all $a, b \in K$. The *differential polynomial ring* $K[t; \delta]$ is the set of polynomials

$$a_0 + a_1t + \cdots + a_nt^n$$

with $a_i \in K$, where addition is defined term-wise and multiplication by

$$ta = at + \delta(a) \quad (a \in K).$$

For $f = a_0 + a_1t + \cdots + a_nt^n$ with $a_n \neq 0$ define $\deg(f) = n$ and $\deg(0) = -\infty$. Then $\deg(fg) = \deg(f) + \deg(g)$. An element $f \in R$ is *irreducible* in R if it is no unit and it has no proper factors, i.e if there do not exist $g, h \in R$ with $\deg(g), \deg(h) < \deg(f)$ such that $f = gh$.

$R = K[t; \delta]$ is a left and right principal ideal domain (a *PID*) and there is a right division algorithm in R : for all $g, f \in R$, $g \neq 0$, there exist unique $r, q \in R$ with $\deg(r) < \deg(f)$, such that

$$g = qf + r.$$

There is also a left division algorithm in R [10, p. 3 and Prop. 1.1.14]. (Our terminology is the one used by Petit [12]; Jacobson's is vice versa.)

Two non-zero elements $f, g \in R$ are called *similar* ($f \sim g$) if and only if there exist $h, q, u \in R$ such that

$$1 = hf + qg \text{ and } u'f = gu$$

for some $u' \in R$ if and only if $R/Rf = R/Rg$ [10, p. 11]. Obviously, $f \sim g$ implies that $\deg(f) = \deg(g)$.

1.3. The characteristic $p > 0$ case. Let K be a field of characteristic p and $R = K[t; \delta]$, then

$$(t - b)^p = t^p - V_p(b), \quad V_p(b) = b^p + \delta^{p-1}(b), \quad (t - b)^{p^e} = t^{p^e} - V_{p^e}(b)$$

for all $b \in K$ with $V_{p^e}(b) = V_p^e(b) = V_p(\dots(V_p(b))\dots)$ [10, p. 17ff]. For any p -polynomial

$$f(t) = a_0t^{p^e} + a_1t^{p^{e-1}} + \cdots + a_et + d \in D[t; \delta]$$

we thus have

$$f(t) - f(t - b) = a_0V_{p^e}(b) + a_1V_{p^{e-1}}(b) + \cdots + a_eb$$

for all $b \in K$ and define

$$V_f(b) = a_0V_{p^e}(b) + a_1V_{p^{e-1}}(b) + \cdots + a_eb.$$

1.4. Nonassociative algebras obtained from differential polynomial rings. Let K be a field and $f \in R = K[t; \delta]$ of degree m . Let $\text{mod}_r f$ denote the remainder of right division by f . Define $F = \text{Cent}(\delta) = \{a \in K \mid \delta(a) = 0\}$.

Definition 1. (cf. [12, (7)]) The vector space

$$R_m = \{g \in K[t; \delta] \mid \deg(g) < m\}$$

together with the multiplication

$$g \circ h = gh \text{ mod}_r f$$

is a unital nonassociative algebra $S_f = (R_m, \circ)$ over

$$F_0 = \{a \in K \mid ah = ha \text{ for all } h \in S_f\}.$$

F_0 is a subfield of K [12, (7)] and it is easy to check that $F_0 = \text{Cent}(\delta)$. The algebra S_f is also denoted by R/Rf [12, 14] if we want to make clear which ring R is involved in the construction. In the following, we call the algebras S_f *Petit algebras* and denote their multiplication simply by juxtaposition. Without loss of generality, we may assume that f is monic, since $S_f \cong S_g$ for all $g = af$ with $a \in K^\times$.

Using left division by f and the remainder $\text{mod}_l f$ of left division by f instead, we can define the multiplication for another unital nonassociative algebra on R_m over F , called ${}_f S$ or R/fR . We will only consider the Petit algebras S_f , since every algebra ${}_f S$ is the opposite algebra of some Petit algebra (cf. [12, (1)]).

Right multiplication with $0 \neq g \in S_f$ is given by $R_g : S_f \rightarrow S_f, h \mapsto hg$, and is a left K -module endomorphism. Left multiplication $L_g : S_f \rightarrow S_f, h \mapsto gh$ is an F -module endomorphism [12], and if we view S_f as a right module over $\text{Nuc}_r(S_f)$, a right $\text{Nuc}_r(S_f)$ -module endomorphism.

Clearly S_f has no zero divisors if and only if R_g and L_g are injective.

Theorem 1. (cf. [12, (2), p. 13-03, (5), (6), (7), (9), (14)]) *Let $f \in R = K[t; \delta]$.*

(i) *If S_f is not associative then $\text{Nuc}_l(S_f) = \text{Nuc}_m(S_f) = K$ and*

$$\text{Nuc}_r(S_f) = \{g \in R_m \mid fg \in Rf\}.$$

The right nucleus of S_f is Amitsur's invariant ring of f .

(ii) *The powers of t are associative if and only if $t^m t = t t^m$ if and only if $t \in \text{Nuc}_r(S_f)$ if and only if $ft \in Rf$.*

(iii) *If f is irreducible then $\text{Nuc}_r(S_f)$ is an associative division algebra.*

(iv) *Let $f \in R$ be irreducible and S_f a finite-dimensional F -vector space or free of finite rank as a right $\text{Nuc}_r(S_f)$ -module. Then S_f is a division algebra.*

Conversely, if S_f is a division algebra then f is irreducible.

(v) *S_f is associative if and only if f is a two-sided element. In that case, S_f is the usual quotient algebra $K[t; \delta]/(f)$.*

(vi) *f is irreducible if and only if S_f is a right division algebra over F (i.e., each non-zero element in S_f has a left inverse: there is $z \in S_f$ such that $zh = 1$), if and only if S_f has no zero divisors.*

Recall that a polynomial $f \in R$ is *bounded* if there exists $0 \neq f^* \in R$, such that $Rf^* = f^*R$ is the largest two-sided ideal of R contained in Rf .

If $f \in R$ is bounded then f is irreducible if and only if $\text{Nuc}_r(S_f)$ has no zero divisors if and only if $\text{Nuc}_r(S_f)$ is an associative division algebra (cf. [7, Proposition 4] which sums up classical results from [9]). [4, Theorem 4] yields:

Theorem 2. *Let $f \in R$ be irreducible. Then f is bounded if and only if S_f is free of finite rank as a $\text{Nuc}_r(S_f)$ -module. In this case, S_f is a division algebra.*

Proof. The first part of the statement is [4, Theorem 4]. Since f irreducible, S_f is a right division algebra and L_h is injective for all $h \in S_f$, $h \neq 0$, as observed in [12, Section 2., (7)]. The second part then follows from the fact that S_f is free of finite rank as a $\text{Nuc}_r(S_f)$ -module, which means the injective $\text{Nuc}_r(S_f)$ -linear map L_h is also surjective. \square

R has finite rank over its center if and only if K is of finite rank over $C_t = \{a \in K \mid at = ta\}$ if and only if all polynomials of R are bounded and if for all f of degree non-zero, $\deg(f^*)/\deg(f)$ is bounded in \mathbb{Q} (f^* being the bound of f) [5, Theorem IV]. Since here $C_t = \text{Const}(\delta) = F$, we conclude:

Proposition 3. *Assume that one of the following equivalent conditions hold:*

- (i) R has finite rank over its center;
- (ii) K/F is a finite field extension.

Then every $f \in R = K[t; \delta]$ is bounded. In particular, if f is irreducible then S_f is a division algebra.

We will assume throughout the paper that $f \in K[t; \delta]$ has $\deg(f) = m \geq 2$ (if f has degree $m = 1$ then $S_f \cong K$) and that $\delta \neq 0$. Without loss of generality, we could only look at monic f , but will do so only when explicitly mentioned.

2. NONASSOCIATIVE ALGEBRAS WHOSE RIGHT NUCLEUS IS A CENTRAL SIMPLE ALGEBRA

We use the terminology from [2] with the only exception that that in our definition of $K[t; \delta]$, we look at polynomials with the coefficients taken on the left, not on the right-hand-side as in [2]. All results, however, work analogously in this case.

Let K be a field of characteristic 0 and δ a derivation of K with $F = \text{Const}(\delta)$, such that K/F is a transcendental extension. Let $f \in R = K[t; \delta]$. The finite-dimensional associative F -algebra $\text{Nuc}_r(S_f)$ is called the *invariant ring* of f by Amitsur [2, p. 260], in modern literature it is also referred to as the *eigenspace* of f .

A differential polynomial $f \in K[t; \delta]$ of degree m is called an *A-polynomial* if there is some $\tilde{f} \in K[t; \delta]$ of degree n such that the resultant $f \times \tilde{f}$ is similar to e_{mn} , the characteristic polynomial of the linear transformation corresponding to the zero $mn \times mn$ matrix [2, p. 263].

The results of Amitsur's paper tell us when $\text{Nuc}_r(S_f)$ is a central simple algebra:

Theorem 4. [2, Lemma 17, 18, 19, Theorem 17, Corollary, Lemma 22] *Let $f, g \in K[t; \delta]$ with $\deg(f) = m \geq 2$ and $\deg(g) = n \geq 2$.*

- (i) $\text{Nuc}_r(S_f)$ has dimension m^2 if and only if f is an *A-polynomial*.
- (ii) If f is an *A-polynomial* then $\text{Nuc}_r(S_f)$ is a central simple algebra of degree m which is

split by K .

(iii) If f and g are A -polynomials then so is $h = f \times g$ and

$$\text{Nuc}_r(S_h) = \text{Nuc}_r(S_f) \otimes_F \text{Nuc}_r(S_g).$$

(iv) If f and g are A -polynomials then

$$\text{Nuc}_r(S_f) \cong \text{Nuc}_r(S_g)$$

if and only if $f \sim g(t+a) \sim g(t) \times t+a$ for some $a \in K$. In particular,

$$S_f \cong S_g \text{ implies that } f \sim g(t+a) \sim g(t) \times t+a$$

for some $a \in K$.

(v) Suppose f is an A -polynomial. Then

$$\text{Nuc}_r(S_f) \cong \text{Mat}_m(F)$$

if and only if one of the following holds:

- $f \sim e_m \times t + c$ for some $c \in K$;
- f decomposes into irreducible factors and at least one factor is linear of the form $t + c$ for some $c \in K$ (then $f \sim e_m \times t + c$).

In particular, then the irreducible factors of f are all similar to $t + c$.

Let L/K be a field extension such that δ extends to L . Then $L[t; \delta]$ is an Ore extension of $K[t; \delta]$ and the constant field $F = \text{Const}(\delta|_K)$ of $\delta = \delta|_K$ is contained in the constant field $C = \text{Const}(\delta)$. If $L = K \cdot C$ is the composite field of K and C , we say L is a *constant extension* of K . It is clear that for $f \in K[t; \delta]$,

$$\text{Nuc}_r(K[t; \delta]/K[t; \delta]f) \subset \text{Nuc}_r(L[t; \delta]/L[t; \delta]f).$$

Theorem 5. Let $f \in K[t; \delta]$ of degree m .

(i) If f is an A -polynomial then $f \in L[t; \delta]$ is an A -polynomial and

$$\text{Nuc}_r(L[t; \delta]/L[t; \delta]f) \cong \text{Nuc}_r(K[t; \delta]/K[t; \delta]f) \otimes_F C.$$

(ii) Suppose $B = \text{Nuc}_r(S_f)$ is a central simple algebra of degree m over F with $f \in K[t; \delta]$. Then C splits B if and only if f has a left or right root in L , i.e. $f = (t-a)g(t) \in L[t; \delta]$ or $f = g(t)(t-a) \in L[t; \delta]$.

In particular, then $S_f \otimes_F C$ has right nucleus $\text{Mat}_m(C)$.

This follows from [2, Theorem 20] and [2, Corollary, p. 270].

Remark 6. Since every automorphism of a nonassociative algebra maps the right nucleus onto itself, for every A -polynomial f which is not two-sided, each $H \in \text{Aut}_F(S_f)$ satisfies $H|_B \in \text{Aut}_F(B)$ when restricted to the central simple algebra $B = \text{Nuc}(S_f)$, thus when restricted to its right nucleus B , is an inner automorphism on B . By an analogous argument, also $H|_K \in \text{Aut}_F(K)$.

3. ALGEBRAS WHOSE RIGHT NUCLEUS IS SPLIT BY A TRANSCENDENTAL EXTENSION

Let F be a field of characteristic 0.

Theorem 7. [2, Lemma 20] (i) *Every central simple algebra B of degree m over F which is split by a transcendental extension K/F is isomorphic to $\text{Nuc}_r(S_f)$ for some $f \in K[t; \delta]$ of degree m and a suitable δ with $F = \text{Const}(\delta)$. The differential polynomial f is an A -polynomial.*

(ii) *Every central division algebra D of degree m over F is isomorphic to $\text{Nuc}_r(S_f)$ for some $f \in K[t; \delta]$ of degree m and a suitable δ with $F = \text{Const}(\delta)$.*

Note that (ii) follows from (i), since for every central division algebra D over F , there is a transcendental field extension K of F in which F is algebraically closed and which splits D ([2, p. 245] or [3]), and we can always find a derivation δ on K with $F = \text{Const}(\delta)$.

As an immediate consequence of Theorem 7 and Remark 6, we now get the following results:

Theorem 8. *For every central simple algebra B of degree m over F which is split by a transcendental field extension K/F , there is a derivation δ on K with field of constants F and a differential polynomial $f \in K[t; \delta]$ of degree m , such that*

$$S_f = K[t; \delta]/K[t; \delta]f$$

is an infinite-dimensional algebra over F with right nucleus B and left and middle nucleus K . Every automorphism $H \in \text{Aut}_F(S_f)$ extends an inner automorphism of B .

We conclude from [2, p. 246]:

Corollary 9. *For every central division algebra D of degree m over F , there exists a transcendental field extension K/F in which F is algebraically closed, a derivation δ on K with field of constants F , and a differential polynomial $f \in K[t; \delta]$ of degree m , such that*

$$S_f = K[t; \delta]/K[t; \delta]f$$

is an infinite-dimensional algebra over F with right nucleus D , and left and middle nucleus K . K splits D and every automorphism $H \in \text{Aut}_F(S_f)$ extends an inner automorphism of D .

The fact that D is a division algebra does not imply that f is irreducible, so S_f might not be a right division algebra.

Corollary 10. *If the differential polynomial f in Corollary 9 is irreducible, then S_f is an infinite-dimensional right division algebra over F and therefore does not have zero divisors.*

If f is an irreducible A -polynomial, it is not bounded by Theorem 2.

Example 11. Suppose $F = \mathbb{R}$. The only central division algebra over \mathbb{R} is $D = (-1, -1)_{\mathbb{R}}$. There exists a transcendental extension K of \mathbb{R} in which \mathbb{R} is algebraically closed, such that K splits D , and a derivation δ on K with $\mathbb{R} = \text{Const}(\delta)$. Thus there is an A -polynomial $f \in K[t; \delta]$ of degree 2, such that

$$S_f = K[t; \delta]/K[t; \delta]f = K \oplus Kt$$

is an infinite-dimensional unital algebra over \mathbb{R} with right nucleus $(-1, -1)_{\mathbb{R}}$, and left and middle nucleus K .

For $B = \text{Mat}_m(\mathbb{R})$ and any transcendental extension K' of \mathbb{R} with a derivation δ on K' where $\mathbb{R} = \text{Const}(\delta)$, there is a reducible A -polynomial $f \in K'[t; \delta]$ of degree m , such that

$$S_f = K'[t; \delta]/K'[t; \delta]f$$

is an infinite-dimensional unital algebra over \mathbb{R} with right nucleus B and left and middle nucleus K' .

4. ALGEBRAS WHOSE RIGHT NUCLEUS IS A p -ALGEBRA

Let now K be a field of characteristic $p > 0$ together with a derivation δ on K . Put $R = K[t; \delta]$ and $F = \text{Const}(\delta)$. There are two cases which can occur: either δ is an algebraic derivation, or δ is transcendental which means $[K : F] = \infty$. We assume that δ is an algebraic derivation of degree p^e with minimum polynomial

$$g(t) = t^{p^e} + c_1 t^{p^{e-1}} + \cdots + c_e t \in F[t]$$

of degree p^e . Then $K = F(u_1, \dots, u_e) = F(u_1) \otimes_F \cdots \otimes_F F(u_e)$ with $u_i^p = a_i \in F$ for all $i \in \{1, \dots, e\}$, and $[K : F] = p^e$, that is K is a finite purely inseparable field extension of exponent one and $K^p \subset F \subset K$. The center $C(R)$ of R is $F[z]$ with $z = g(t) - d_0$, $d_0 \in F$ and the two-sided elements in R have the form $uh(t)$ with $u \in K^\times$, $h(t) \in C(R)$.

Recall that a central simple algebra $B = \text{Mat}_r(D)$ over a field F of characteristic p is a p -algebra if it has index p^n , equivalently, if its exponent is a power of p [10, p. 154].

Note that for $f(t) = g(t) - d \in F[t]$ (so $f(t)$ is two-sided in this case),

$$(K, \delta, d) = K[t; \delta]/K[t; \delta]f(t)$$

is an associative central simple F -algebra called a *differential extension of K* and treated in [10, p. 23]. K is a maximal subfield of (K, δ, d) .

Theorem 12. [2, Lemma 20'] *Let B be a p -algebra of degree m over F which is split by a purely inseparable extension K of exponent one (i.e., has exponent p), such that $m \leq [K : F]$. Then*

$$B \cong \text{Nuc}_r(S_f)$$

for some $f \in K[t; \delta]$ of degree m and a suitable δ with $F = \text{Const}(\delta)$.

We start by looking at the case that $m = [K : F] = p^e$ and immediately obtain (i) and (ii) in the following result on p -algebras by employing only Theorem 1 (v) from Petit [12] and Amitsur's Theorem 12 (only the fact that then B is cyclic uses Hood's Main Theorem [8, Main Theorem]):

Theorem 13. *Let B be a p -algebra of degree m over F which is split by a purely inseparable field extension K of exponent one with $m = [K : F]$.*

(i) *There is an algebraic derivation δ on K of degree m with minimum polynomial $g(t)$ such that the center of $K[t; \delta]$ is $F[z]$ with $z = g(t) - d_0$, $d_0 \in F$, and*

$$B = (K, \delta, d_0)$$

with $f(t) = g(t) - d_0$. B is a cyclic algebra.

(ii) $F[t]/(f)$ is a subfield of B of degree p^e over F if and only if f is irreducible in $F[t]$.

(iii) $f \in K[t; \delta]$ is irreducible if and only if B is a division algebra.

(iv) $B \cong \text{Mat}_{p^e}(F)$ if and only if there is $b \in K$ such that

$$d_0 = V_g(b) = V_{p^e}(b) + c_1 V_{p^{e-1}}(b) + \cdots + c_e b.$$

Proof. (i) If $m = [K : F]$ then there is a differential polynomial $f \in K[t; \delta]$ of degree m and a suitable δ such that $B \cong \text{Nuc}_r(S_f)$ by Theorem 12. Here B is an associative subalgebra of S_f of dimension m^2 and S_f has dimension m^2 as well. Therefore $S_f = B$ is associative and $f \in K[t; \delta]$ must be a two-sided differential polynomial of degree m , i.e. $B = K[t; \delta]/(f)$ is a quotient algebra (Theorem 1 (v)). Without loss of generality we may assume f is monic. Thus $f \in C(R)$ and since f has degree $m = p^e$, we obtain that $f(t) = g(t) - d_0$ and so $B = (K, \delta, d_0)$. K is a purely inseparable field extension of F which is an (even maximal) subfield of B splitting B , therefore B is cyclic [8, Main Theorem].

(ii) Since here $f(t) \in F[t]$, we know that $F[t]/(f)$ is a subfield of B of degree p^e over F if and only if f is irreducible in $F[t]$.

(iii) is [7, Proposition 4] and (iv) is a consequence from (i) together with Theorem [10, Theorem 1.3.27]. \square

Remark 14. Let us briefly put the previous result into context:

(i) Let A be a central simple p -algebra of degree p^n over F . It is a well known classical result that A is cyclic over F if and only if A has a subfield K such that K is a purely inseparable extension of F and K is a splitting field for A (this is [8, Main Theorem], which removed Albert's restriction that K be simple from [1, Theorem (7.27)]).

(ii) Mammone characterized the central simple algebras split by a purely inseparable field extension K of exponent one in [11]: in particular, if B is a central simple algebra over F of degree $m = p^e$ containing K where $[K : F] = m$, then B is a *differential crossed product*, that means B contains a K -basis of the form $\{z_1^{i_1} \cdots z_n^{i_n} \mid 0 \leq i_k \leq p - 1\}$ satisfying a kind of commutativity law with elements of K which involves a set of n F -derivations of K . The algebra B then yields elements $b_i = z_i^p$ and $u_{ij} = z_i z_j - z_j z_i$ in K . Conversely, given sets $B = \{b_i \mid i = 1, \dots, n\}$ and $U = \{u_{ij} \mid i, j = 1, \dots, n\}$ satisfying certain relations involving F -derivations of K , then (U, B) arises from such a differential crossed product.

In case $m < [K : F] = p^e$ we obtain a nonassociative algebra of dimension mp^e containing B as right nucleus:

Theorem 15. *Let B be a p -algebra of degree m over F which is split by a purely inseparable extension K of exponent one such that $m < [K : F]$.*

(i) *There is an algebraic derivation δ and a differential polynomial $f \in K[t; \delta]$ of degree m such that*

$$S_f = K[t; \delta]/K[t; \delta]f$$

is an algebra over F of dimension mp^e with right nucleus B , left and middle nucleus K , and nucleus $\text{Nuc}(S_f) = B \cap K$ an intermediate field of K/F , unequal to K .

(ii) *f is irreducible if and only if B is a division algebra, if and only if S_f is a division*

algebra.

(iii) Every automorphism $H \in \text{Aut}_F(S_f)$ extends an inner automorphism of B and an automorphism of K .

Proof. (i) The existence of a suitable f follows from Theorem 12 and the statements on the left and middle nuclei from Theorem 1. Since f is not two-sided, K is not contained in the right nucleus of S_f , i.e. not contained in B [13, Theorem 9]. Thus $\text{Nuc}(S_f) = B \cap K$ is properly contained in K , so that it is an intermediate field of the field extension K/F .

(ii) By Proposition 3 and Theorem rethm:structure, f is irreducible if and only if B is a division algebra, if and only if S_f is a division algebra.

(iii) An automorphism of S_f extends both an inner automorphism of B and an automorphism of K by Remark 6. \square

Corollary 16. *Let D be a division p -algebra of degree m over F which is split by a purely inseparable extension K of exponent one such that $m < [K : F]$. Then there is an irreducible polynomial $f \in K[t; \delta]$ of degree m such that S_f is a division algebra over F of dimension mp^e with right nucleus D , left and middle nucleus K , and nucleus $D \cap K$ an intermediate field of K/F , unequal to K .*

The fact that f is irreducible in Corollary 16 follows from Proposition 3. Note that every division p -algebra over F split by K has degree $m \leq [K : F]$, so that Theorem 13 (iii) and Corollary 16 cover all possible cases for a division p -algebra.

We could ask for the algebra S_f of smallest possible dimension which contains a given central simple algebra B as a right nucleus. This is equivalent to asking for a purely inseparable extension K of exponent one splitting B of smallest possible degree $[K : F] = p^e$ satisfying $m < [K : F]$, which in turn is connected to the question how many cyclic algebras are needed when saying that B is similar to a product of cyclic algebras of degree p in the Brauer group $Br(F)$.

Theorem 17. *Let B be a p -algebra over F of degree m , index $d = p^n$ and exponent p , such that $m = r^2 p^n < p^{d-1}$. Then there is a purely inseparable extension K of exponent one with $[K : F] = p^{d-1}$, and a differential polynomial $f \in K[t; \delta]$ of degree m such that*

$$S_f = K[t; \delta]/K[t; \delta]f$$

is an algebra over F of dimension mp^{d-1} with right nucleus B and the properties listed in Theorem 15

Proof. Let B be a p -algebra of index p^n and exponent p . Then there is a purely inseparable field extension K/F of exponent one with $K = F(u_1, \dots, u_{d-1})$, $u_i^p = a_i \in F$, and $[K : F] = p^{d-1}$, which splits B [6, Theorem 1.1.]. We have $m = r^2 d = r^2 p^n$ for some $r \geq 1$.

We need $m = r^2 p^n \leq [K : F] = p^{d-1}$ to be able to apply Theorem 12. By Theorem 12 this implies that $B \cong \text{Nuc}_r(S_f)$ for some $f \in K[t; \delta]$ of degree m and a suitable δ with $F = \text{Const}(\delta)$. Since each $f \in K[t; \delta]$ is bounded by Proposition 3, $B = \text{Nuc}_r(S_f)$ is a division algebra if and only if f is irreducible [7, Proposition 4], if and only if S_f is a division algebra. \square

We obtain that for a division algebra D , the smallest possible dimension l of a division algebra S_f containing D as right nucleus satisfies $m^2 < l = mp^e \leq mp^{m-1}$:

Corollary 18. *Let D be a division p -algebra of degree m and exponent p over F . Then there is a purely inseparable extension K of exponent one with $[K : F] = p^{m-1}$, and an irreducible differential polynomial $f \in K[t; \delta]$ of degree m such that*

$$S_f = K[t; \delta]/K[t; \delta]f$$

is a division algebra over F of dimension mp^{m-1} with right nucleus D and the properties listed in Theorem 15.

Proof. There is a purely inseparable field extension $K = F(u_1, \dots, u_{m-1})$ of exponent one, $u_i^p = a_i \in F$, and $[K : F] = p^{m-1}$, which splits D [6, Theorem 1.1.].

We need $m = p^n \leq [K : F] = p^{m-1}$ to be able to apply Theorem 12. This holds for all prime p and $n \geq 1$ as it is equivalent to $n \leq p^n - 1$, i.e. to $n + 1 \leq p^n$, which is true for all prime p and $n \geq 1$. Therefore there is a purely inseparable field extension K/F of exponent one with $m \leq [K : F] = p^{m-1}$ which splits D . By Theorem 12 this implies that $B \cong \text{Nuc}_r(S_f)$ for some $f \in K[t; \delta]$ of degree m and a suitable δ with $F = \text{Const}(\delta)$. Since D is a division algebra and f bounded, f is irreducible and S_f is a division algebra. \square

REFERENCES

- [1] A. A. Albert, "Structure of algebras." Revised printing, Amer. Math. Soc., Providence, R.I., 1961.
- [2] A. S. Amitsur, *Differential Polynomials and Division Algebras*. Annals of Mathematics, Vol. 59 (2) (1954) 245-278.
- [3] A. S. Amitsur, *Generic splitting fields of central simple algebras*. Ann. of Math. 62 (2) (1955), 8-43.
- [4] J. Carcanague, *Idéaux bilatres d'un anneau de polynomes non commutatifs sur un corps*. J. Algebra (18) 1971, 1-18.
- [5] J. Carcanague, *Quelques résultats sur les anneaux de Ore*. C. R. Acad. Sci. Paris Sr. A-B 269 (1969), A749-A752.
- [6] M. Florence, *On the symbol length of p -algebras*. Compositio Mathematica 149 (8) (2013), 1353-1363.
- [7] J. Gómez-Torrecillas, *Basic module theory over non-commutative rings with computational aspects of operator algebras. With an appendix by V. Levandovskyy*. Lecture Notes in Comput. Sci. 8372, Algebraic and algorithmic aspects of differential and integral operators, Springer, Heidelberg (2014) 23-82.
- [8] J. M. Hood, *Central simple p -algebras with purely inseparable subfields*. J. Algebra 17 (1971) 299-301.
- [9] N. Jacobson, *The Theory of Rings*, AMS, Providence, RI, 1943
- [10] N. Jacobson, "Finite-dimensional division algebras over fields." Springer Verlag, Berlin-Heidelberg-New York, 1996.
- [11] P. Mammone, *Remarques sur les produits croiss différentiels*. Acad. Roy. Belg. Bull. Cl. Sci. (5) 68 (1982), no. 10, 651-664.
- [12] J.-C. Petit, *Sur certains quasi-corps généralisant un type d'anneau-quotient*. Séminaire Dubriel. Algèbre et théorie des nombres 20 (1966 - 67), 1-18.
- [13] S. Pumplün, *Nonassociative differential extensions over fields of characteristic p* . Preprint 2016.
- [14] J.-C. Petit, *Sur les quasi-corps distributives à base momogène*. C. R. Acad. Sc. Paris 266 (1968), Série A, 402-404.
- [15] R. D. Schafer, "An Introduction to Nonassociative Algebras." Dover Publ., Inc., New York, 1995.
E-mail address: susanne.pumpluen@nottingham.ac.uk

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF NOTTINGHAM, UNIVERSITY PARK, NOTTINGHAM NG7 2RD, UNITED KINGDOM