

ALBERT ALGEBRAS OVER \mathbb{Z} AND OTHER RINGS

SKIP GARIBALDI, HOLGER P. PETERSSON, AND MICHEL L. RACINE

ABSTRACT. Albert algebras, a specific kind of Jordan algebra, are naturally distinguished objects among commutative non-associative algebras and also arise naturally in the context of simple affine group schemes of type F_4 , E_6 , or E_7 . We study these objects over an arbitrary base ring R , with particular attention to the case $R = \mathbb{Z}$. We prove in this generality results previously in the literature in the special case where R is a field of characteristic different from 2 and 3.

CONTENTS

1. Introduction	1
2. Notation	3
3. Background on polynomial laws	3
4. Background on composition algebras	6
5. Background on Jordan algebras	8
6. Cubic Jordan algebras	9
7. Albert algebras are Freudenthal algebras are Jordan algebras	12
8. The ideal structure of Freudenthal algebras	14
9. Groups of type F_4 and C_3	15
10. Generic minimal polynomial of a Freudenthal algebra	17
11. Basic classification results for Albert algebras	19
12. Isotopy	20
13. Classification of Albert algebras over \mathbb{Z}	23
14. Groups of type E_6	24
15. Freudenthal triple systems	27
16. Groups of type E_7	29
17. Exceptional groups over \mathbb{Z}	31
References	32

1. INTRODUCTION

Albert algebras, a specific kind of Jordan algebra, are naturally distinguished objects among commutative non-associative algebras and also arise naturally in the context of simple affine group schemes of type F_4 , E_6 , or E_7 . We study these objects over an arbitrary base ring R , with particular attention to the case $R = \mathbb{Z}$. We prove in this generality results previously in the literature in the special case where R is a field of characteristic different from 2 and 3.

1991 *Mathematics Subject Classification*. Primary 17C40; Secondary 17C30, 20G41.

Why Albert algebras? In the setting of semisimple algebraic groups over a field, a standard technique for computing with elements of a group — especially an anisotropic group — is to interpret the group in terms of automorphisms of some algebraic structure, such as viewing an adjoint group of type B_n as the special orthogonal group of a quadratic form of dimension $2n + 1$, or an adjoint group of type A_n as the automorphism group of an Azumaya algebra of rank $(n + 1)^2$. This approach can be seen in many references, from [Wei60], through [KMRT98] and [Con]. In this vein, Albert algebras appear as a natural tool for computations related to F_4 , E_6 , and E_7 groups, as we do below.

In the setting of nonassociative algebras, Albert algebras arise naturally. Among commutative not-necessarily-associative algebras under additional mild hypotheses (the field has characteristic $\neq 2, 3, 5$ and the algebra is metrized), every algebra satisfying a polynomial identity of degree ≤ 4 is a Jordan algebra, see [ChG, Prop. A.8]. Jordan algebras have an analogue of the Wedderburn-Artin theory for associative algebras [Jac68, p. 201, Cor. 2], and one finds that all the simple Jordan algebras are closely related to associative algebras (more precisely, “are special”) except for one kind, the Albert algebras, see for example [Jac68, p. 210, Th. 11] or [MZ88].

Our contribution. In the setting of nonassociative algebras, we prove a classification of Albert algebras over \mathbb{Z} (Theorem 13.3), which was viewed as an open question in the context of nonassociative algebra; here we see that it is equivalent to the classification of groups of type F_4 , which was known, see [Con] which leverages [Gro96] and [EG96]. We also prove new results about ideals in Albert algebras (Theorem 8.3) and about isotopy of Albert algebras over local rings (Theorem 12.5). We have not seen Lemma 14.1 in the literature, even in the case of a base field of characteristic different from 2 and 3.

In the setting of affine group schemes, we present Albert algebras in a streamlined way in Definition 7.1. Note that this definition is in the context of what was formerly called a “quadratic” Jordan algebra — because instead of a bilinear multiplication one has a quadratic map, the U -operator — and that it makes sense whether or not 2 is invertible in the base ring. Applying this definition here allows one to replace, in some proofs, “global” computations over \mathbb{Z} as one finds in [Con] with “local” computations over an algebraically closed field that exist in several places in the literature (see, for example, the proof of Lemma 9.1). We also interpret a clever computation in [EG96] as an example of a general mechanism known as isotopy, see Definition 13.1. Our classification of groups of type E_7 over \mathbb{Z} in Proposition 17.1 uses general techniques to reduce the problem to computations over \mathbb{R} .

Comparison with other works. The survey [Pet19] also considers Albert algebras over rings. It asserts that $\mathbf{Aut}(J)$ is a smooth group scheme of type F_4 for J an Albert algebra, saying that the proof is similar to the analogous result for octonion algebras and groups of type G_2 in [LPR08]. We give a different and complete proof here, see Lemma 9.1.

The definition of Freudenthal algebra in [Pet19] is different from here, but the two definitions are essentially equivalent, see Remark 7.5.

A recent article by Alsaody, [Als], gives several interesting examples about Albert algebras over rings, especially concerning isotopy, compare §12 here. That paper relies on the assertion about $\mathbf{Aut}(J)$ already mentioned.

Changing our viewpoint away from nonassociative algebras and towards group schemes, this note owes various debts to [Con].

2. NOTATION

Rings, by definition, have a 1. We put $\mathbb{Z}\text{-alg}$ for the category of commutative rings, where \mathbb{Z} is an initial object. For any $R \in \mathbb{Z}\text{-alg}$, we put $R\text{-alg}$ for the category of pairs (S, f) with $S \in \mathbb{Z}\text{-alg}$ and $f: R \rightarrow S$, i.e., the coslice category $R \downarrow \mathbb{Z}\text{-alg}$. Below, R will typically denote an element of $\mathbb{Z}\text{-alg}$. (The interested reader is invited to mentally replace R by a base scheme X , $R\text{-alg}$ with the category of schemes over X , finitely generated projective R -modules with vector bundles over X , etc., thereby translating results below into a language closer to that in [CalF].)

We write $\text{Mat}_n(R)$ for the ring of n -by- n matrices with entries from R ; its invertible elements form the group $\text{GL}_n(R)$. We write $\langle \alpha_1, \dots, \alpha_n \rangle \in \text{Mat}_n(R)$ for the diagonal matrix whose (i, i) -entry is α_i .

Suppose now that \mathbf{G} is a finitely presented group scheme over R . For each fppf $S \in R\text{-alg}$, we write $H^1(S/R, \mathbf{G})$ for the collection of \mathbf{G} -torsors over R trivialized by S , see for example [Gir71], [Wat79], or [CalF, §2.2]. It does not depend on the choice of structure homomorphism $R \rightarrow S$ [Gir71, Rem. III.3.6.5]. The subcategory of fppf elements of $R\text{-alg}$ has a small skeleton, so the union

$$H^1(R, \mathbf{G}) := \bigcup_{\text{fppf } S \in R\text{-alg}} H^1(S/R, \mathbf{G})$$

is a set. It is the non-abelian fppf cohomology of \mathbf{G} . In case \mathbf{G} is smooth, it agrees with étale H^1 . If additionally R is a field, then it agrees with the non-abelian Galois cohomology defined in, for example, [Ser02].

Unimodular elements. Let M be an R -module. An element $m \in M$ is said to be *unimodular* if Rm is a free R -module of rank 1 and a direct summand of M , equivalently, if there is some $\lambda \in M^*$ (the dual of M) such that $\lambda(m) = 1$. When M is finitely generated projective, this is equivalent to: $m \otimes 1$ is not zero in $M \otimes F$ for every field $F \in R\text{-alg}$, see for example [Loo06, 0.3].

If $m \in M$ is unimodular, then so is $m \otimes 1 \in M \otimes S$ for every $S \in R\text{-alg}$. In the opposite direction, if M is finitely generated projective, S is a cover of R (i.e., $\text{Spec } S \rightarrow \text{Spec } R$ is surjective), and $m \otimes 1$ is unimodular in $M \otimes S$, it follows that m is unimodular as an element of M .

3. BACKGROUND ON POLYNOMIAL LAWS

We may identify an R -module M with a functor $\mathbf{W}(M)$ from $R\text{-alg}$ to the category of sets defined via $S \mapsto M \otimes S$. For R -modules M, N , a polynomial law (in the sense of [Rob63]) $f: \mathbf{W}(M) \rightarrow \mathbf{W}(N)$ is a morphism of functors, i.e., a collection of set maps $f_S: M \otimes S \rightarrow N \otimes S$ varying functorially with S . We put $\mathcal{P}_R(M, N)$ for the collection of polynomial laws $\mathbf{W}(M) \rightarrow \mathbf{W}(N)$, and omit the subscript R when it is understood.

A polynomial law is *homogeneous of degree* $d \geq 0$ if $f_S(sx) = s^d f_S(x)$ for every $S \in R\text{-alg}$, $s \in S$, and $x \in M \otimes S$, see [Rob63, p. 226]. A *form of degree* d on M is a polynomial law $\mathbf{W}(M) \rightarrow \mathbf{W}(R)$ that is homogeneous of degree d . The forms of degree 0 are constants, i.e., given by an element of R . Those of degree 1 are R -linear maps $M \rightarrow R$. Those of degree 2 are commonly known as quadratic forms on M . We put $\mathcal{P}_R^d(M, N)$ for the submodule of $\mathcal{P}_R(M, N)$ of polynomial laws that are homogeneous of degree d .

It is often useful to argue that a polynomial law f is zero, which a priori means checking a condition for all $S \in R\text{-alg}$. However, it suffices to verify that $f_T = 0$ for every *local*

ring $T \in R\text{-alg}$. Indeed, for $m \in M \otimes S$, $f_S(m) = 0$ in $N \otimes S$ if and only if $f_S(m) \otimes 1 = f_{S_{\mathfrak{m}}}(m \otimes 1) = 0$ in $N \otimes S_{\mathfrak{m}}$ for every maximal ideal \mathfrak{m} of S .

Lemma 3.1. *Let M be a finitely generated projective R -module, and suppose $f \in \mathcal{P}(M, R)$ is such that $f(0) = 0$. If $m \in M$ has $f(m) \in R^\times$, then m is unimodular.*

Proof. If m is not unimodular, then there is a field $F \in R\text{-alg}$ such that $m \otimes 1 = 0$ in $M \otimes F$, and $f(m \otimes 1) = 0$, whence $f(m)$ belongs to the kernel of $R \rightarrow F$, a contradiction. \square

Directional derivatives. For $f \in \mathcal{P}(M, N)$, $v \in M$, and t an indeterminate $n \geq 0$, we define a polynomial law $\nabla_v^n f$ as follows. For $S \in R\text{-alg}$ and $x \in M \otimes S$, $f_{S[t]}(x + v \otimes t)$ is an element of $N \otimes S[t]$, and we define $\nabla_v^n f_S(x) \in N \otimes S$ to be the coefficient of t^n . This defines a polynomial law called the n -th directional derivative $\nabla_v^n f$ of f in the direction v . One finds that $\nabla_v^0 f = f$ regardless of v . We abbreviate $\nabla_v f := \nabla_v^1 f$; it is linear in v .

If f is homogeneous of degree d and $0 \leq n \leq d$, then $\nabla_v^n f(x)$ is homogeneous of degree $d - n$ in x and degree n in v . The symmetry implicit in the definition of the directional derivative gives $\nabla_v^n f(x) = \nabla_x^{d-n} f(v)$ for $x \in M$.

Lemma 3.2. *Suppose M, N are R -modules and A is a unital associative R -algebra and $g \in \mathcal{P}(M, A)$ is a polynomial law such that there is an element $m \in M$ such that $g(m) \in A$ is invertible. If $f \in \mathcal{P}^d(M, N)$ satisfies*

$$g_S(x) \in A_S^\times \Rightarrow f_S(x) = 0$$

for all $S \in R\text{-alg}$ and $x \in M \otimes S$, then f is identically zero.

Proof. Since the hypotheses are stable under base change, it suffices to show that $f(v) = 0$ for all $v \in M$. Replacing g by $L \circ g \in \mathcal{P}(M, A)$, where $L \in \text{End}_R(A)$ is multiplication in A on the left by the inverse of $g(m)$, we may assume $g(m) = 1_A$. Set $S := R[\varepsilon]/(\varepsilon^{d+1})$. For $v \in M$, the element

$$g_S(m + \varepsilon v) = 1_A + \sum_{n=1}^d \varepsilon^n \nabla_v^n g(m)$$

is invertible in A_S , so by hypothesis,

$$0 = f_S(m + \varepsilon v) = \sum_{n=0}^d \varepsilon^n \nabla_v^n f(m).$$

Focusing on the coefficient of ε^d in that equation gives

$$0 = \nabla_v^d f(m) = \nabla_m^0 f(v) = f(v),$$

as required. \square

The module of polynomial laws. In the following, we write $S^n M$ for the n -th symmetric power of M , i.e., the R -module $\otimes^n M$ modulo the submodule generated by elements $x - \sigma(x)$ for $x \in \otimes^n M$ and σ a permutation of the n factors.

Lemma 3.3. *Let M and N be finitely generated projective R -modules. Then for each $d \geq 0$:*

- (1) $\mathcal{P}^d(M, N)$ is a finitely generated projective R -module.
- (2) If $T \in R\text{-alg}$ is faithfully flat, the natural map $\mathcal{P}_R^d(M, N) \otimes T \rightarrow \mathcal{P}_T^d(M \otimes T, N \otimes T)$ is an isomorphism.
- (3) The natural map $S^d(M^*) \otimes N \rightarrow \mathcal{P}^d(M, N)$ is an isomorphism.
- (4) The natural map $\mathcal{P}^d(M, R) \otimes N \rightarrow \mathcal{P}^d(M, N)$ is an isomorphism.

Proof. $\mathcal{P}_R^d(M, N)$ is naturally isomorphic to $\text{Hom}_R(\Gamma_d(M), N)$ by [Rob63, Th. IV.1], where $\Gamma_d(M)$ denotes the module of degree d divided powers on M . Then $\mathcal{P}_R^d(M, N) \otimes T \cong \text{Hom}_R(\Gamma_d(M), N) \otimes T$, which in turn is $\text{Hom}_T(\Gamma_d(M) \otimes T, N \otimes T)$ because T is faithfully flat [KO74, p. 33, Prop. II.2.5]. Now $\Gamma_d(M) \otimes T \cong \Gamma_d(M \otimes T)$ by [Bour, §IV.5, Exercise 7], completing the proof of (2).

(3): If M and N are free modules, then the map is an isomorphism by [Rob63, p. 232]. If M and N have constant rank, then there is a faithfully flat $T \in R\text{-alg}$ such that $M \otimes T$ and $N \otimes T$ are free. Since (3) holds over T by the free case, (2) and faithfully flat descent give that (3) holds. In the general case, since M and N are finitely generated projective, we may write $R = \prod_{i=0}^n R_i$ for some n such that $M = \bigoplus M_i$ and $N = \bigoplus N_i$ with each M_i, N_i an R_i -module of finite constant rank. Then $\mathcal{P}^d(M, N) = \bigoplus \mathcal{P}^d(M_i, N_i)$ and $S^d(M^*) \otimes N = \bigoplus (S^d(M_i^*) \otimes N_i)$ and the claim follows by the constant rank case.

(4) follows trivially from (3). For (1), note that M^* is finitely generated projective, so so is $S^d(M^*)$ and also the tensor product $S^d(M^*) \otimes N$. Applying (3) gives the claim. \square

One can create new polynomial laws from old by twisting by a line bundle.

Lemma 3.4. *Let M and N be finitely generated projective R -modules. Then for every $d \geq 0$ and every line bundle L , we have:*

- (1) *There is a natural isomorphism $\mathcal{P}^d(M, N) \otimes (L^*)^{\otimes d} \rightarrow \mathcal{P}^d(M \otimes L, N)$.*
- (2) *There is a natural isomorphism $\mathcal{P}^d(M, N) \cong \mathcal{P}^d(M \otimes L, N \otimes L^{\otimes d})$.*

Proof. For (1), since L^* is a line bundle, the natural map $(L^*)^{\otimes d} \rightarrow S^d(L^*)$ is an isomorphism because it is so after faithfully flat base change. Since $S^d(M^*) \otimes S^d(L^*)$ is naturally identified with $S^d((M \otimes L)^*)$, combining Lemma 3.3(3),(4) then gives (1).

For (2), there are isomorphisms $\mathcal{P}^d(M \otimes L, N \otimes L^{\otimes d}) \xrightarrow{\sim} \mathcal{P}^d(M, N) \otimes (L^*)^{\otimes d} \otimes L^{\otimes d}$ by (1) and Lemma 3.3(4). Since $L^{\otimes d} \otimes (L^*)^{\otimes d} \cong R$, the claim follows. \square

Example 3.5. Suppose L is a line bundle and there is an isomorphism $h: L^{\otimes d} \rightarrow R$ for some $d \geq 1$. Such pairs $[L, h]$ are called (approximately) d -trivialized line bundles in [CalF, §2.4.3] and were studied in the case $d = 2$ in [Knu91], where they are called *discriminant modules*. Applying h to identify $N \otimes L^{\otimes d} \xrightarrow{\sim} N$ in Lemma 3.4(2) gives a construction that takes $f \in \mathcal{P}^d(M, N)$ and gives an element of $\mathcal{P}^d(M \otimes L, N)$, which we denote by $[L, h] \cdot (M, f)$.

For example, for each $\alpha \in R^\times$, define $\langle \alpha \rangle$ to be $[L, h]$ as in the preceding paragraph, where $L = R$ and h is defined by $h(\ell_1 \otimes \cdots \otimes \ell_d) = \alpha \prod \ell_i$. Clearly, $\langle \alpha \beta^d \rangle \cong \langle \alpha \rangle$ for all $\alpha, \beta \in R^\times$. Applying the construction in the previous paragraph, we find $\langle \alpha \rangle \cdot (M, f) \cong (M, \alpha f)$.

Every $[L, h]$ with $L = R$ is necessarily isomorphic to $\langle \alpha \rangle$ for some $\alpha \in R^\times$. In particular, if $\text{Pic}(R)$ has no d -torsion elements other than zero — e.g., if R is a semilocal ring or a UFD [Sta18, tags 0BCH, 02M9] — then each $[L, h]$ is isomorphic to $\langle \alpha \rangle$ for some α .

The group scheme μ_d of d -th roots of unity is the automorphism group of each $[L, h]$, where μ_d acts by multiplication on L . The group $H^1(R, \mu_d)$ classifies pairs (L, h) .

We say that homogeneous polynomial laws related by the isomorphism in Lemma 3.4(2) are *projectively similar*, imitating the language from [AuBB, §1.2] for the case of quadratic forms ($d = 2$). (This relationship was called “lax-similarity” in [BC].) We say that homogeneous degree d laws f and $[L, h] \cdot f$ for $[L, h] \in H^1(R, \mu_d)$ as in the preceding example are *similar*. If $\text{Pic}(R)$ has no d -torsion elements other than zero, the two notions coincide.

For $f \in \mathcal{P}^d(M, N)$, we define $\mathbf{Aut}(f)$ to be the subgroup of $\mathbf{GL}(M)$ consisting of elements g such that $fg = f$ as polynomial laws. In case M and N are finitely generated projective, so is $\mathcal{P}^d(M, N)$, whence the functor $\mathbf{Aut}(f)$ from $R\text{-alg}$ to groups defined by $\mathbf{Aut}(f)(T) = \mathbf{Aut}(f_T)$ is a closed sub-group-scheme of $\mathbf{GL}(M)$.

Lemma 3.6. *Let f and f' be homogeneous polynomial laws on finitely generated projective modules. If f and f' are projectively similar, then their automorphism groups are isomorphic.*

Proof. By hypothesis, $f \in \mathcal{P}^d(M, N)$ and $f' \in \mathcal{P}^d(M \otimes L, N \otimes L^{\otimes d})$ for some modules M and N ; line bundle L ; and $d \geq 0$. The group scheme $\mathbf{Aut}(f)$ is the closed subgroup-scheme of $\mathbf{GL}(M)$ stabilizing the element f in $\mathbf{S}^d(M^*) \otimes N$. Now, any element of $\mathbf{GL}(M)$ acts on $\mathbf{S}^d((M \otimes L)^*) \otimes (N \otimes L^{\otimes d})$ by defining it to act as the identity on L . In this way, we find a homomorphism $\mathbf{Aut}(f) \rightarrow \mathbf{Aut}(f')$. Viewing M as $(M \otimes L) \otimes L^*$ and N as $(N \otimes L^{\otimes d}) \otimes (L^*)^{\otimes d}$, and repeating this construction, we find an inverse mapping $\mathbf{Aut}(f') \rightarrow \mathbf{Aut}(f)$. \square

4. BACKGROUND ON COMPOSITION ALGEBRAS

A not-necessarily-associative R -algebra C is an R -module with an R -linear map $C \otimes_R C \rightarrow C$, which we view as a multiplication and write as juxtaposition. Such a C is *unital* if it has an element $1_C \in C$ such that $1_C c = c 1_C = c$ for all $c \in C$. See e.g. [Sch94]. A *composition R -algebra* as in [Pet93] is such a C that is finitely generated projective as an R -module, is unital, and has a quadratic form $n_C : C \rightarrow R$ that allows composition (that is, such that $n_C(xy) = n_C(x)n_C(y)$ for all $x, y \in C$), satisfies $n_C(1_C) = 1$, and whose bilinearization defined by $n_C(x, y) := n_C(x+y) - n_C(x) - n_C(y)$ gives an isomorphism $C \rightarrow C^*$ via $x \mapsto n_C(x, \cdot)$. We say that a symmetric bilinear form with this property is *regular*. The quadratic form n_C (which is unique by Proposition 4.5 below) is called the *norm* of C .

Remark 4.1. In the definition above, one can swap the condition $n_C(1_C) = 1$ with the requirement that the rank of C is nowhere zero.

We put $\mathrm{Tr}_C(x) := n_C(x, 1_C)$, a linear map $C \rightarrow R$, called the *trace* of C . Trivially, $\mathrm{Tr}_C(1_C) = 2$. Lemma 3.1 gives that 1_C is unimodular, so we may identify R with $R1_C$, and C is a faithful R -module. The unimodularity of 1_C is equivalent to the existence of some $\lambda \in C^*$ such that $\lambda(1_C) = 1$, i.e., some $x \in C$ such that $\mathrm{Tr}_C(x) = 1$, whence $\mathrm{Tr}_C : C \rightarrow R$ is surjective.

The class of composition algebras is stable under base change. That is, if C is a composition R -algebra with norm n_C , then for every $S \in R\text{-alg}$, $C \otimes S$ is a composition S -algebra with norm $n_C \otimes S$. The following two results are essentially well known [Pet93, 1.2–1.4]. For convenience, we include their proof.

Lemma 4.2 (“Cayley-Hamilton”). *Let C be a composition algebra with norm n_C and define Tr_C as above. Then*

$$x^2 - \mathrm{Tr}_C(x)x + n_C(x)1_C = 0$$

for all $x \in C$.

Proof. Linearizing the composition law $n_C(xy) = n_C(x)n_C(y)$, we find

$$(4.3) \quad n_C(xy, x) = n_C(x) \mathrm{Tr}_C(y) \quad \text{and}$$

$$(4.4) \quad n_C(xy, wz) + n_C(wy, xz) = n_C(x, w)n_C(y, z)$$

for all $x, y, z, w \in C$. Setting $z = x$ and $w = 1_C$ in (4.4), we find:

$$n_C(xy, x) + n_C(y, x^2) = \text{Tr}_C(x)n_C(x, y).$$

Combining these with (4.3), we find:

$$n_C(x^2 - \text{Tr}_C(x)x + n_C(x)1_C, y) = 0 \quad \text{for all } x, y \in C.$$

Since the bilinear form n_C is regular, the claim follows. \square

A priori, a composition algebra is a unital algebra together with a quadratic form, the norm. The next result shows that this data is redundant.

Proposition 4.5. *If C is a composition algebra, then the norm n_C is uniquely determined by the algebra structure of C .*

Proof. Let $n' : C \rightarrow R$ be any quadratic form making C a composition algebra and write Tr' for the corresponding trace $\text{Tr}'(x) := n'(x + 1_C) - n'(x) - n'(1_C)$. Then $\lambda := \text{Tr}_C - \text{Tr}'$ (resp., $q := n_C - n'$) is a linear (resp., quadratic) form on C and the Cayley-Hamilton property yields

$$(4.6) \quad \lambda(x)x = q(x)1_C \quad \text{for all } x \in C.$$

We aim to prove that $q = 0$. Because 1_C is unimodular, it suffices to prove $\lambda = 0$. This can be checked locally, so we may assume that R is local and in particular $C = R1_C \oplus M$ for a free module M . Now, $\text{Tr}_C(1_C) = 2 = \text{Tr}'(1_C)$, so $\lambda(1_C) = 0$. For $m \in M$ a basis vector, $\lambda(m)m$ belongs to $M \cap R1_C$ by (4.6), so it is zero, whence $\lambda(m) = 0$, proving the claim. \square

Corollary 4.7. *Let C be a unital R -algebra. If there is a faithfully flat $S \in R\text{-alg}$ such that $C \otimes S$ is a composition S -algebra, then C is a composition algebra over R .*

Proof. Because the norm $n_{C \otimes S}$ of $C \otimes S$ is uniquely determined by the algebra structure, one obtains by faithfully flat descent a quadratic form $n_C : C \rightarrow R$ such that $n_C \otimes S = n_{C \otimes S}$. Because $n_{C \otimes S}$ satisfies the properties required to make $C \otimes S$ a composition algebra and S is faithfully flat over R , it follows that the same properties hold for n_C . \square

The following facts are standard, see for example [Knu91, §V.7]: Composition algebras are alternative algebras. The map $\bar{\cdot} : C \rightarrow C$ defined by $\bar{x} := \text{Tr}_C(x)1_C - x$ is an involution, i.e., an R -linear anti-automorphism of period 2.

Composition algebras of constant rank. In case R is connected, a composition R -algebra has rank 2^e for $e \in \{0, 1, 2, 3\}$ [Knu91, p. 206, Th. V.7.1.6]. Therefore, specifying a composition R -algebra C is equivalent to writing

$$(4.8) \quad R = \prod_{e=0}^3 R_e \quad \text{and} \quad C = \prod_{e=0}^3 C_e,$$

where C_e is a composition R_e -algebra of constant rank 2^e .

If C is a composition algebra of rank 1, then since 1_C is unimodular, C is equal to R . The bilinear form $n_C(\cdot, \cdot)$ gives an isomorphism $C \rightarrow C^*$ and $n_C(1_C, \alpha 1_C) = 2\alpha$, we deduce that 2 is invertible in R . Conversely, if 2 is invertible, then R is a composition algebra by setting $n_C(\alpha) = \alpha^2$.

A composition algebra whose rank is 2 is not just an associative and commutative ring, it is an étale algebra [Knu91, p. 43, Th. I.7.3.6]. Conversely, every rank 2 étale algebra is a composition algebra. Among rank 2 étale algebras, there is a distinguished one, $R \times R$, which is said to be split.

A composition algebra whose rank is 4 is associative and is an Azumaya algebra, commonly known as a *quaternion algebra*. (Note that our notion of quaternion algebra is more restrictive than the one in the books [Knu91, see p. 43] and [Voi21].) Among quaternion R -algebras, there is a distinguished one, the 2-by-2 matrices $\text{Mat}_2(R)$, which is said to be split.

A composition algebras whose rank is 8 is known as an *octonion algebra*. Among octonion R -algebras, there is a distinguished one that is said to be split, called the Zorn vector matrices and denoted $\text{Zor}(R)$, see [LPR08, 4.2]. As a module, we view it as $\begin{pmatrix} R & R^3 \\ R^3 & R \end{pmatrix}$ with multiplication

$$\begin{pmatrix} \alpha_1 & u \\ x & \alpha_2 \end{pmatrix} \begin{pmatrix} \beta_1 & v \\ y & \beta_2 \end{pmatrix} = \begin{pmatrix} \alpha_1\beta_1 - u^\top y & \alpha_1v + \beta_2u + x \times y \\ \beta_1x + \alpha_2y + u \times v & -x^\top v + \alpha_2\beta_2 \end{pmatrix}$$

where \times is the ordinary cross product on R^3 . The quadratic form is

$$n_{\text{Zor}(R)} \begin{pmatrix} \alpha_1 & u \\ x & \alpha_2 \end{pmatrix} = \alpha_1\alpha_2 + u^\top x.$$

One says that a composition R -algebra C is *split* if, when we write R and C as in (4.8), C_e is isomorphic to the split composition R_e -algebra for $e \geq 1$.

Example 4.9. The real octonions \mathbb{O} are a composition \mathbb{R} -algebra with basis $1_{\mathbb{O}}, e_1, e_2, \dots, e_7$ which is orthonormal with respect to the quadratic form $n_{\mathbb{O}}$ with multiplication table

$$e_r^2 = -1 \quad \text{and} \quad e_r e_{r+1} e_{r+3} = -1$$

for all r with subscripts taken modulo 7, and the displayed triple product is associative.

The \mathbb{Z} -sublattice \mathcal{O} of \mathbb{O} spanned by $1_{\mathbb{O}}$, the e_r , and

$$\begin{aligned} h_1 &= (1 + e_1 + e_2 + e_4)/2, & h_2 &= (1 + e_1 + e_3 + e_7)/2, \\ h_3 &= (1 + e_1 + e_5 + e_6)/2, & \text{and } h_4 &= (e_1 + e_2 + e_3 + e_5)/2 \end{aligned}$$

is a composition \mathbb{Z} -algebra. It is a maximal order in $\mathcal{O} \otimes \mathbb{Q}$, and all such are conjugate under the automorphism group of $\mathcal{O} \otimes \mathbb{Q}$. (As a consequence, there is some choice in the way one presents this algebra. We have followed [EG96].) As a subring of \mathbb{O} , it has no zero divisors. For more on this, see [Dic23, §19], [Cox], [ConwS, §9], or [Con, §5].

5. BACKGROUND ON JORDAN ALGEBRAS

Para-quadratic and Jordan algebras. A (unital) *para-quadratic algebra* over a ring R is an R -module J together with a quadratic map $U: R \rightarrow \text{End}_R(J)$ — i.e., U is an element of $\mathcal{P}^2(R, \text{End}_R(J))$ — called the *U-operator*, and a distinguished element $1_J \in J$, such that $U_{1_J} = \text{Id}_J$. As a notational convenience, we define a linear map $J \otimes J \otimes J \rightarrow J$ denoted $x \otimes y \otimes z \mapsto \{xyz\}$ via

$$(5.1) \quad \{xyz\} := (U_{x+z} - U_x - U_z)y.$$

Evidently, $\{xyz\} = \{zyx\}$ for all $x, y, z \in J$.

A para-quadratic R -algebra J is a *Jordan R -algebra* if the identities

$$(5.2) \quad U_{U_x y} = U_x U_y U_x \quad \text{and} \quad U_x \{y x z\} = \{(U_x y) z x\}$$

hold for all $x, y, z \in J \otimes S$ for all $S \in R\text{-alg}$. (Alternatively, one can define a Jordan R -algebra entirely in terms of identities concerning elements of J , avoiding the “for all $S \in R\text{-alg}$ ”, at the cost of requiring a longer list of identities, see [McC66, §1].) Note that if J is a Jordan R -algebra, then $J \otimes T$ is a Jordan T -algebra for every $T \in R\text{-alg}$ (“Jordan algebras are closed under base change”). If J is a para-quadratic algebra and $J \otimes T$ is Jordan for some faithfully flat $T \in R\text{-alg}$, then J is Jordan.

For x in a Jordan algebra J and $n \geq 0$, we define the n -th power x^n via

$$(5.3) \quad x^0 := 1_J, \quad x^1 := x, \quad x^n = U_x x^{n-2} \text{ for } n \geq 2.$$

An element $x \in J$ is *invertible with inverse* y if $U_x y = x$ and $U_x y^2 = 1$ [McC66, §5]. It turns out that x is invertible if and only if U_x is invertible if and only if 1 is in the image of U_x ; when these hold, the inverse of x is $y = U_x^{-1}x$, which we denote by x^{-1} . It follows from (5.2) that $x, y \in J$ are both invertible if and only if $U_x y$ is invertible, and in this case $(U_x y)^{-1} = U_{x^{-1}y^{-1}}$.

Example 5.4. Let A be an associative and unital R -algebra. Define $U_x y := xyx$ for $x, y \in A$. Then $\{xyz\} = xyz + zyx$ and A endowed with this U -operator is a Jordan algebra denoted by A^+ . Note that for $x \in A$ and $n \geq 0$, the n -th powers of x in A and A^+ are the same.

Relations with other kinds of algebras. Suppose for this paragraph that 2 is invertible in R . Given a para-quadratic algebra J as in the preceding paragraph, one can define a commutative (bilinear) product \bullet on J via

$$(5.5) \quad x \bullet y := \frac{1}{2}\{x1_J y\} \quad \text{for } x, y \in J.$$

(In the case where J is constructed from an associative algebra as in Example 5.4, one finds that $x \bullet y = \frac{1}{2}(xy + yx)$. If additionally the associative algebra is commutative, \bullet equals the product in that associative algebra.) If J is Jordan, then \bullet satisfies

$$(5.6) \quad (x \bullet y) \bullet (x \bullet x) = x \bullet (y \bullet (x \bullet x)),$$

which is the axiom classically called the ‘‘Jordan identity’’.

In the opposite direction, given an R -module J with a commutative product \bullet with identity element 1_J , we obtain a para-quadratic algebra by setting

$$(5.7) \quad U_x y := 2x \bullet (x \bullet y) - (x \bullet x) \bullet y \quad \text{for } x, y \in J.$$

If the original product satisfied the Jordan identity, then the para-quadratic algebra so obtained satisfies (5.2), i.e., is a Jordan algebra in our sense, see for example [McC04, p. 202].

Definition 5.8 (hermitian matrix algebras). Let C be a composition R -algebra and $\Gamma = \langle \gamma_1, \gamma_2, \gamma_3 \rangle \in \text{GL}_3(R)$. We define $\text{Her}_3(C, \Gamma)$ to be the R -submodule of $\text{Mat}_3(C)$ consisting of elements fixed by the involution $x \mapsto \Gamma^{-1} \bar{x}^T \Gamma$ and with diagonal entries in R . Note that, as an R -module, $\text{Her}_3(C, \Gamma)$ is a sum of 3 copies of C and 3 copies of R , so it is finitely generated projective.

In the special case where 2 is invertible in R , one can define a multiplication \bullet on $\text{Her}_3(C, \Gamma)$ via $x \bullet y := \frac{1}{2}(xy + yx)$, where juxtaposition denotes the usual product of matrices in $\text{Mat}_3(C)$. It satisfies the Jordan identity [Jac68, p. 61, Cor.], and therefore the U -operator defined via (5.7) makes $\text{Her}_3(C, \Gamma)$ into a Jordan algebra.

6. CUBIC JORDAN ALGEBRAS

In this section, we define cubic Jordan algebras and the closely related notion of cubic norm structure. They provide a useful alternative language for computation.

Definition 6.1. Following [McC69] (see [PR86, p. 212] for the terminology), we define a *cubic norm R -structure* as a quadruple $\mathbf{M} = (M, 1_{\mathbf{M}}, \sharp, N_{\mathbf{M}})$ consisting of an R -module M , a distinguished element $1_{\mathbf{M}} \in M$ (the *base point*), a quadratic map $\sharp: M \rightarrow M$, $x \mapsto x^\sharp$ (the *adjoint*), with (symmetric bilinear) polarization $x \times y := (x + y)^\sharp - x^\sharp - y^\sharp$, a

cubic form $N_{\mathbf{M}}: M \rightarrow R$ (the *norm*) such that the following axioms are fulfilled. Defining a bilinear form $T_{\mathbf{M}}: M \times M \rightarrow R$ by

$$(6.2) \quad T_{\mathbf{M}}(x, y) := (\nabla_x N_{\mathbf{M}})(1_{\mathbf{M}})(\nabla_y N_{\mathbf{M}})(1_{\mathbf{M}}) - (\nabla_x \nabla_y N_{\mathbf{M}})(1_{\mathbf{M}})$$

(the *bilinear trace*), which is symmetric since the directional derivatives ∇_x, ∇_y commute [Rob63, p. 241, Prop. II.5], and a linear form $\text{Tr}_{\mathbf{M}}: M \rightarrow R$ by

$$(6.3) \quad \text{Tr}_{\mathbf{M}}(x) := T_{\mathbf{M}}(x, 1_{\mathbf{M}})$$

(the *linear trace*), the identities

$$(6.4) \quad 1_{\mathbf{M}}^{\sharp} = 1_{\mathbf{M}}, \quad N_{\mathbf{M}}(1_{\mathbf{M}}) = 1,$$

$$(6.5) \quad 1_{\mathbf{M}} \times x = \text{Tr}_{\mathbf{M}}(x)1_{\mathbf{M}} - x, \quad (\nabla_y N_{\mathbf{M}})(x) = T_{\mathbf{M}}(x^{\sharp}, y), \quad x^{\sharp\sharp} = N_{\mathbf{M}}(x)x$$

hold in all scalar extensions $M \otimes S, S \in R\text{-alg}$.

For such a cubic norm structure \mathbf{M} , we then define a U -operator by

$$(6.6) \quad U_x y := T_{\mathbf{M}}(x, y)x - x^{\sharp} \times y,$$

which together with $1_{\mathbf{M}}$ converts the R -module M into a Jordan R -algebra $J = J(\mathbf{M})$ [McC69, Th. 1]. In the sequel, we rarely distinguish carefully between the cubic norm structure \mathbf{M} and the Jordan algebra $J(\mathbf{M})$. By abuse of notation, we write $1_J = 1_{\mathbf{M}}$, $N_J = N_{\mathbf{M}}$, $T_J = T_{\mathbf{M}}$, and $\text{Tr}_J := \text{Tr}_{\mathbf{M}}$ if there is no danger of confusion, even though, in general, J does not determine \mathbf{M} uniquely [PR86, p. 216].

A Jordan R -algebra J is said to be *cubic* if there exists a cubic norm R -structure \mathbf{M} as above such that (i) $J = J(\mathbf{M})$ and (ii) $J = M$ is a finitely generated projective R -module. With the quadratic form $S_J: M \rightarrow R$ defined by $S_J(x) := \text{Tr}_J(x^{\sharp})$ for $x \in J$ (the *quadratic trace*), the cubic Jordan algebra J satisfies the identities

$$(6.7) \quad (U_x y)^{\sharp} = U_{x^{\sharp}} y^{\sharp}, \quad N_J(U_x y)U_x y = N_J(x)^2 N_J(y)U_x y,$$

$$(6.8) \quad U_x x^{\sharp} = N_J(x)x, \quad U_x(x^{\sharp})^2 = N_J(x)^2 1_J,$$

$$(6.9) \quad x^{\sharp} = x^2 - \text{Tr}_J(x)x + S_J(x)1_J = 0, \quad \text{and}$$

$$(6.10)$$

$$x^3 - \text{Tr}_J(x)x^2 + S_J(x)x - N_J(x)1_J = 0 = x^4 - \text{Tr}_J(x)x^3 + S_J(x)x^2 - N_J(x)x$$

for all $x \in J$. For (6.7)–(6.9) and the first equation of (6.10), see [McC69, p. 499], while the second equation of (6.10) follows from the first, (6.8), and (6.9) via $x^4 = U_x x^2 = U_x x^{\sharp} + \text{Tr}_J(x)U_x x - S_J(x)U_x 1_J = \text{Tr}_J(x)x^3 - S_J(x)x^2 + N_J(x)x$.

Remark 6.11. Note that the second equality of (6.10) derives from the first through formal multiplication by x . But, due to the para-quadratic character of Jordan algebras, this is not a legitimate operation unless 2 is invertible in R . In fact, cubic Jordan algebras exist that contain elements x satisfying $x^2 = 0 \neq x^3$ [Jac69, 1.31–1.32].

Lemma 6.12. *Let J be a cubic Jordan R -algebra and $x, y \in J$.*

- (1) *x is invertible in J if and only if $N_J(x)$ is invertible in R . In this case*

$$x^{-1} = N_J(x)^{-1} x^{\sharp} \quad \text{and} \quad N_J(x^{-1}) = N_J(x)^{-1}.$$

- (2) *Invertible elements of J are unimodular.*

- (3) *$N_J(U_x y) = N_J(x)^2 N_J(y)$ and $N_J(x^2) = N_J(x)^2 = N_J(x^{\sharp})$.*

Proof. (1): If $N_J(x)$ is invertible in R , then (6.8) shows that so is x , with inverse $x^{-1} = N_J(x)^{-1}x^\sharp$. Conversely, assume x is invertible in J . Then $y := (x^{-1})^2$ satisfies $U_x y = 1_J$, and (6.7) yields $1_J = N_J(U_x y)U_x y = N_J(x)^2 N_J(y)1_J$, hence

$$N_J(x)^2 N_J(y) = 1$$

since 1_J is unimodular by Lemma 3.1 and (6.4). Thus $N_J(x) \in R^\times$. Before proving the final formula of (1), we deal with (2), (3).

(2) follows immediately from Lemma 3.1 combined with the first part of (1).

(3): Applying Lemma 3.2 to the polynomial law $g: J \times J \rightarrow \text{End}_R(J)$ defined by $g(x, y) := U_{U_x y}$ in all scalar extensions, we may assume that $U_x y$ is invertible. By (2), therefore, $U_x y$ is unimodular, and the first equality follows from (6.7). The second equality follows from the first for $y = 1_J$, while in the third equality we may again assume that x is invertible, hence unimodular. Then (6.8) combines with the first equality to imply $N_J(x)^4 = N_J(N_J(x)x) = N_J(U_x x^\sharp) = N_J(x)^2 N_J(x^\sharp)$, as desired.

Now the second equality of (1) follows from the first and (3) via

$$N_J(x^{-1}) = N_J(x)^{-3} N_J(x^\sharp) = N_J(x)^{-1}. \quad \square$$

Without the assumption that J is finitely generated projective as an R -module, Lemma 6.12 would be false [PR85, Th. 10].

Example 6.13. We endow the R -module $M := \text{Her}_3(C, \Gamma)$ from Definition 5.8 with a cubic norm R -structure $\mathbf{M} = (M, 1_{\mathbf{M}}, \sharp, N_{\mathbf{M}})$, where $1_{\mathbf{M}}$ is the 3-by-3 identity matrix. An element of $x \in \text{Her}_3(C, \Gamma)$ may be written as

$$x = \begin{pmatrix} \alpha_1 & \gamma_2 c_3 & \gamma_3 \bar{c}_2 \\ \gamma_1 \bar{c}_3 & \alpha_2 & \gamma_3 c_1 \\ \gamma_1 c_2 & \gamma_2 \bar{c}_1 & \alpha_3 \end{pmatrix}$$

for $\alpha_i \in R$ and $c_i \in C$. Because three of the entries are determined by symmetry, we may denote such an element by

$$(6.14) \quad x := \sum_{i=1}^3 (\alpha_i \varepsilon_i + \delta_i^\Gamma(c_i)),$$

where ε_i has a 1 in the (i, i) entry and zeros elsewhere, and $\delta_i^\Gamma(c)$ has $\gamma_{i+2}c$ in the $(i+1, i+2)$ entry — where the symbols $i+1$ and $i+2$ are taken modulo 3 — and zeros in the other entries not determined by symmetry. In the literature on Jordan algebras, one finds the notation $c[(i+1)(i+2)]$ for what we denote $\delta_i(c)$.

We define the adjoint \sharp by

$$x^\sharp := \sum_{i=1}^3 \left((\alpha_{i+1} \alpha_{i+2} - \gamma_{i+1} \gamma_{i+2} n_C(c_i)) \varepsilon_i + \delta_i^\Gamma(-\alpha_i c_i + \gamma_i \overline{c_{i+1} c_{i+2}}) \right)$$

with indices mod 3, and the norm $N_{\mathbf{M}}$ by

$$(6.15) \quad N_{\mathbf{M}}(x) := \alpha_1 \alpha_2 \alpha_3 - \sum_{i=1}^3 \gamma_{i+1} \gamma_{i+2} \alpha_i n_C(c_i) + \gamma_1 \gamma_2 \gamma_3 \text{Tr}_C(c_1 c_2 c_3)$$

in all scalar extensions, where the last summand on the right of (6.15) is unambiguous since $\text{Tr}_C((c_1 c_2) c_3) = \text{Tr}_C(c_1 (c_2 c_3))$ [McC85, Th. 3.5]. By [McC69, Th. 3], \mathbf{M} is indeed a cubic norm structure. The corresponding cubic Jordan algebra will again be denoted by $J := \text{Her}_3(C, \Gamma) := J(\mathbf{M})$.

(In case 2 is invertible in R , the commutative product \bullet on $\text{Her}_3(C, \Gamma)$ defined from the U -operator by (5.5) equals the product $x \bullet y := \frac{1}{2}(xy + yx)$ from Definition 5.8. In order to see this, it suffices to note that the square of $x \in \text{Her}_3(C, \Gamma)$ as defined in (5.3) is the same as the square of x in the matrix algebra $\text{Mat}_3(C)$. This in turn follows immediately from (6.9), (6.15), and the definition of the adjoint.)

For x as above and $y = \sum(\beta_i \varepsilon_i + \delta_i^\Gamma(v_i))$, with $\beta_i \in R$, $v_i \in C$, evaluating the bilinear trace at x, y yields

$$(6.16) \quad T_J(x, y) = \sum_{i=1}^3 (\alpha_i \beta_i + \gamma_{i+1} \gamma_{i+2} n_C(u_i, v_i)).$$

Since the bilinear trace n_C is regular, so is T_J .

For the special case where $\Gamma = \text{Id}$, we define $\text{Her}_3(C) := \text{Her}_3(C, \text{Id})$ and write δ_i for δ_i^Γ . It can be useful to write elements of $\text{Her}_3(C)$ as

$$\begin{pmatrix} \alpha_1 & c_3 & \cdot \\ \cdot & \alpha_2 & c_1 \\ c_2 & \cdot & \alpha_3 \end{pmatrix}$$

where \cdot denotes an entry that is omitted because it is determined by symmetry. As an example of the triple product defined from (5.1) and (6.6), we mention that for $x = \sum \alpha_i \varepsilon_i$ diagonal, we have

$$(6.17) \quad \{\delta_i(a) \delta_{i+1}(b) x\} = \delta_{i+2}(\overline{ab}) \alpha_i \quad \text{and} \quad \{\delta_{i+1}(b) \delta_i(a) x\} = \delta_{i+2}(\overline{ab}) \alpha_{i+1}$$

for $i \in 1, 2, 3$ taken mod 3 and $a, b \in C$.

Note that, for the Jordan algebra $\text{Her}_3(C, \Gamma)$ just defined, if we multiply Γ by an element of R^\times or any entry in Γ by the square of an element of R^\times , we obtain an algebra isomorphic to the original. Therefore, replacing Γ by $\langle (\det \Gamma)^{-1} \gamma_1, (\det \Gamma) \gamma_2, (\det \Gamma) \gamma_3 \rangle$ does not change the isomorphism class of $\text{Her}_3(C, \Gamma)$ and we may assume that $\gamma_1 \gamma_2 \gamma_3 = 1$.

Example 6.18. In case $R = \mathbb{R}$, the preceding paragraph shows that it is sufficient to consider two choices for Γ , namely $\langle 1, 1, \pm 1 \rangle$. We compute $T_{\text{Her}_3(C, \Gamma)}$ for each choice of C and Γ . Regular symmetric bilinear forms over \mathbb{R} are classified by their dimension and signature (an integer), so it suffices to specify the signature. If $C = \mathbb{R}, \mathbb{C}, \mathbb{H}$, or \mathbb{O} , the signature of n_C is 2^r for $r = 0, 1, 2, 3$ respectively. By (6.16), T_J has signature $3(1 + 2^r)$ for $J = \text{Her}_3(C)$ and $3 - 2^r$ for $J = \text{Her}_3(C, \langle 1, 1, -1 \rangle)$. For J the split Freudenthal algebra of rank $3(1 + 2^r)$ with $r = 1, 2$, or 3 , the signature of T_J is 3.

Remark 6.19. Alternatively, one could define the Jordan algebra structure on $\text{Her}_3(C, \Gamma)$ for an arbitrary ring R without referring to cubic norm structures as follows. Writing out the formulas for the U -operator from Definition 5.8 in case $R = \mathbb{Q}$, one finds that the formulas do not involve any denominators other than γ_i terms and therefore make sense for any R regardless of whether 2 is invertible. This makes $\text{Her}_3(C, \Gamma)$ a para-quadratic algebra. Because it is a Jordan algebra in case $R = \mathbb{Q}$ as in Definition 5.8, we conclude that $\text{Her}_3(C, \Gamma)$ is a Jordan algebra with no hypothesis on R by extension of identities [Bour, §IV.2.3, Th. 2]. This alternative definition gives the same objects, but is much harder to work with.

7. ALBERT ALGEBRAS ARE FREUDENTHAL ALGEBRAS ARE JORDAN ALGEBRAS

Definition 7.1. A *split Freudenthal R -algebra* is a Jordan algebra $\text{Her}_3(C)$ as in Example 6.13 for some split composition R -algebra C . Because split composition algebras are determined up to isomorphism by their rank function, so are split Freudenthal algebras.

A para-quadratic R -algebra J is a *Freudenthal algebra* if $J \otimes S$ is a split Freudenthal S -algebra for some faithfully flat $S \in R\text{-alg}$. It is immediate that every Freudenthal algebra is a Jordan algebra.

Since every split Freudenthal R -algebra is finitely generated projective as an R -module for every R , the same is true for every Freudenthal R -algebra J [Sta18, Tags 03C4, 05A9], and by the same reasoning we see that the identity element 1_J is unimodular. Because the

rank of a composition algebra takes values in $\{1, 2, 4, 8\}$, the rank of a Freudenthal algebra takes values in $\{6, 9, 15, 27\}$. An *Albert R -algebra* is a Freudenthal R -algebra of rank 27.

Proposition 7.2. *For every composition R -algebra C and every $\Gamma \in \mathrm{GL}_3(R)$, $\mathrm{Her}_3(C, \Gamma)$ is a Freudenthal algebra.*

Proof. Replacing R with R_e as in (4.8), we may assume that C has constant rank. There is a faithfully flat $S \in R\text{-alg}$ such that $C \otimes S$ is a split composition algebra.

Consider $T := S[t_1, t_2, t_3]/(t_1^2 - \gamma_1, t_2^2 - \gamma_2, t_3^2 - \gamma_3)$. It is a free S -module, so faithfully flat. Then $\mathrm{Her}_3(C, \Gamma) \otimes T$ is isomorphic to $\mathrm{Her}_3(C \otimes T)$ as cubic Jordan algebras, and the latter is a split Freudenthal algebra. \square

The Freudenthal algebras $\mathrm{Her}_3(C, \Gamma)$ are said to be *reduced*.

Example 7.3. Let J be a Freudenthal R -algebra. If $x \in J$ has $U_x = \mathrm{Id}_J$, then $x = \zeta 1_J$ for $\zeta \in R$ such that $\zeta^2 = 1$. To see this, first suppose that J is $\mathrm{Her}_3(C)$ for some composition algebra C and write $x = \sum(\alpha_i \varepsilon_i + \delta_i(c_i))$ for $\alpha_i \in R$ and $c_i \in C$. We find

$$U_x \varepsilon_i = \alpha_i^2 \varepsilon_i + \delta_{i+2}(\alpha_i c_{i+2}) + \cdots$$

for each i , so $\alpha_i^2 = 1$ and $c_{i+2} = 0$ for all i . Then

$$U_x \delta_i(1_C) = \delta_i(\alpha_{i+1} \alpha_{i+2} 1_C).$$

Since 1_C is unimodular, $\alpha_{i+1} \alpha_{i+2} = 1$ for all i , proving the claim for this J .

For general J , let $S \in R\text{-alg}$ be faithfully flat such that $J \otimes S$ is split. Then $x \in J$ maps to an element of $R 1_J \otimes S \subseteq J \otimes S$ and so belongs to $R 1_J \subseteq J$. Since $U_{\zeta 1_J} = \zeta^2 \mathrm{Id}_J$ for $\zeta \in R$, the claim follows.

The following result is well known when R is a field or perhaps a local ring, see for example [Pet19, Prop. 20]. We impose no hypothesis on R .

Proposition 7.4. *Suppose C is a split composition R -algebra of constant rank at least 2, i.e., C is $R \times R$, $\mathrm{Mat}_2(R)$, or $\mathrm{Zor}(R)$. Then $\mathrm{Her}_3(C, \Gamma) \cong \mathrm{Her}_3(C)$ for all Γ .*

Proof. Since n_C is universal, there are invertible $p, q \in C$ such that $\gamma_2 = n_C(q^{-1})$ and $\gamma_3 = n_C(p^{-1})$, so $\gamma_1 = n_C(pq)$. We define $C^{(p,q)}$ to be a not-necessarily associative R -algebra with the same underlying R -module structure and with multiplication $\cdot_{(p,q)}$ defined by

$$x \cdot_{(p,q)} y := (xp)(qy),$$

where the multiplication on the right is the multiplication in C . Certainly $(pq)^{-1}$ is an identity element in $C^{(p,q)}$. The algebra $C^{(p,q)}$ is called an *isotope* of C and is studied in [McC71a], where it is proved to be alternative. One checks that it is a composition algebra with quadratic form $n_{C^{(p,q)}} = n_C(pq)n_C$, see [McC71a, Prop. 5] for a more general statement in case R is a field.

Define $\phi : \mathrm{Her}_3(C^{(p,q)}) \rightarrow \mathrm{Her}_3(C, \Gamma)$ via $\phi(\sum x_i \varepsilon_i + \delta_i(c_i)) = \sum x_i \varepsilon_i + \delta_i^\Gamma(c'_i)$, where

$$c'_1 = (pq)c_1(pq), \quad c'_2 = c_2 p, \quad \text{and} \quad c'_3 = q c_3.$$

It is evidently an isomorphism of R -modules and one checks that it is an isomorphism of Jordan algebras, compare [McC71a, Th. 3]. Therefore, we are reduced to verifying that $C^{(p,q)}$ is split.

If C is associative, then the R -linear map

$$L_{pq} : C^{(p,q)} \rightarrow C \quad \text{such that} \quad L_{pq}(x) = pqx$$

is an isomorphism of R -algebras. So assume $C = \mathrm{Zor}(R)$.

At the beginning, when we chose p and q , we were free to pick $\xi_i, \eta_i \in R^\times$ such that $p = \begin{pmatrix} \xi_1 & 0 \\ 0 & \xi_2 \end{pmatrix}$ and $q = \begin{pmatrix} \eta_1 & 0 \\ 0 & \eta_2 \end{pmatrix}$. Let $A \in \text{Mat}_3(R)$ be any matrix such that $\det A = (\xi_1 \xi_2^2 \eta_1^2 \eta)^{-1}$ and put $B := \xi_2 \eta_1 (A^\sharp)^\top$, where \sharp denotes the classical adjoint. With $\zeta_i := (\xi_i \eta_i)^{-1}$, one checks, using the formula $(Sx) \times (Sy) = (S^\sharp)^\top(x \times y)$ for \times the usual cross product in R^3 , that the assignment

$$\begin{pmatrix} \alpha_1 & u_1 \\ u_2 & \alpha_2 \end{pmatrix} \mapsto \begin{pmatrix} \zeta_1 \alpha_1 & Au_1 \\ Bu_2 & \zeta_2 \alpha_2 \end{pmatrix}$$

defines an isomorphism $C \xrightarrow{\sim} C^{(p,q)}$. \square

Remark 7.5. We have defined Albert algebras by a sort of descent condition in Definition 7.1. In view of Proposition 10.2 below, one could alternatively define an Albert algebra as a cubic Jordan algebra that is projective of rank 27 as an R -module such that $J \otimes F$ is a simple Jordan F -algebra for every field F ; this is the definition used in [Pet19] and [Als], for example. The two notions lead to the same objects.

8. THE IDEAL STRUCTURE OF FREUDENTHAL ALGEBRAS

It is a standard exercise to show that every (two-sided) ideal in the matrix algebra $\text{Mat}_n(R)$ is of the form $\text{Mat}_n(\mathfrak{a})$ for some ideal \mathfrak{a} in R . More generally, every ideal in an Azumaya R -algebra A is of the form $\mathfrak{a}A$ some ideal \mathfrak{a} of R [KO74, p. 95, Cor. III.5.2].

A similar result holds for every octonion R -algebra C : *Every one-sided ideal in C is a two-sided ideal that is stable under the involution on C . The maps $I \mapsto I \cap R$ and $\mathfrak{a}C \mapsto \mathfrak{a}$ are bijections between the set of ideals of C and ideals in R . See [Pet21, §4] for a proof in this generality and the references therein for earlier results of this type going back to [Mah42].*

We now prove a similar result for Freudenthal algebras.

Definition 8.1. An *ideal* in a para-quadratic R -algebra J is the kernel of a homomorphism, i.e., an R -submodule I such that

$$U_I J + U_J I + \{JJI\} = I,$$

where we have written $U_I J$ for the R -span of $U_x y$ with $x \in I$ and $y \in J$. (This is sometimes written with a \subseteq instead of $=$, but the two are equivalent since $U_J I \supseteq U_{1_J} I = I$.) An R -submodule I is an *outer ideal* if

$$(8.2) \quad U_J I + \{JJI\} = I.$$

Here are some observations about outer ideals:

- (1) Every ideal is an outer ideal.
- (2) If 2 is invertible in R , then for every $x \in I$ and $y \in J$, $U_x y = \frac{1}{2}\{xyx\} \in \{JJI\}$, so the notions of ideal and outer ideal coincide.
- (3) For every ideal \mathfrak{a} in R , the R -submodule $\mathfrak{a}J$ is an ideal of J .
- (4) If 1_J is unimodular, then for every outer ideal I of J , $I \cap R1_J$ is an ideal in R , for the trivial reason that I is an R -module.
- (5) If \mathfrak{a} is an ideal in R and 1_J is unimodular, then $\mathfrak{a}1_J = (\mathfrak{a}J) \cap R1_J$. The containment \subseteq is clear. To see the opposite containment, suppose $\alpha 1_J \in \mathfrak{a}J \cap R1_J$ for some $\alpha \in R$ and write $\alpha 1_J = \sum \alpha_i y_i$ with $\alpha_i \in \mathfrak{a}$ and $y_i \in J$. There is some R -linear $\lambda: J \rightarrow R$ such that $\lambda(1_J) = 1$. Then $\alpha = \lambda(\alpha 1_J) = \sum \alpha_i \lambda(y_i)$ is in \mathfrak{a} .

Theorem 8.3. *Let J be a Freudenthal R -algebra. Every outer ideal of J is an ideal. The maps $I \mapsto I \cap R1_J$ and $\mathfrak{a}J \mapsto \mathfrak{a}$ are bijections between the set of outer ideals of J and the set of ideals of R .*

Proof. It suffices to show that the stated maps are bijections, because then observation (3) implies that every outer ideal is of the form $\mathfrak{a}J$ and therefore an ideal. In view of (5) (noting that 1_J is unimodular), it suffices to verify that $(I \cap R1_J)J = I$ for every outer ideal I .

First suppose that $J = \text{Her}_3(C)$ for some composition R -algebra C and write $\mathfrak{a} := I \cap R1_J$. The Peirce projections relative to the diagonal frame of J , i.e., U_{ε_i} and $x \mapsto \{\varepsilon_j x \varepsilon_l\}$ for $i, j, l = 1, 2, 3$ [McC66, p. 1074] stabilize I , and we find

$$I = \sum_i (I \cap R\varepsilon_i) + (I \cap \delta_i(C)).$$

Set $B := \{c \in C \mid \delta_1(c) \in I\}$. We claim that B is an ideal in C . Note that $U_{\delta_1(1_C)}\delta_1(b) = \delta_1(\bar{b})$, so B is stable under the involution.

We leverage (6.17). Repeatedly applying this with $a = 1_C$ and using that B is stable under the involution, we conclude that $\delta_i(B) = I \cap \delta_i(C)$ for all i . For $c \in C$ and $b \in B$, I contains $\{1_J \delta_2(\bar{c}) \delta_1(\bar{b})\} = \delta_3(cb)$, so $cB \subseteq B$, i.e., B is an ideal in C and therefore $B = \mathfrak{a}C$ for some ideal \mathfrak{a} of R .

For $c \in C$, I contains $\{\delta_i(1_C)\varepsilon_{i+1}\delta_i(\mathfrak{a}c)\} = \text{Tr}_C(\mathfrak{a}c)\varepsilon_{i+2}$. Since Tr_C is surjective, $\mathfrak{a}\varepsilon_j \subseteq I$ for all j .

In the other direction, if $\alpha_i \varepsilon_i \in I$, then so is

$$\{\delta_{i+1}(1_C)1_J(\alpha_i \varepsilon_i)\} = \delta_{i+1}(\alpha_i 1_C).$$

It follows that $I \cap R\varepsilon_i = \mathfrak{a}R$ for all i and in particular, $I \cap R1_J = \mathfrak{a}R$ and $I = \mathfrak{a}J$.

We now treat the general case. Suppose I is an outer ideal in a Freudenthal R -algebra J . There is a faithfully flat $S \in R\text{-alg}$ such that $J \otimes S$ is a split Freudenthal algebra. We have

$$((I \cap R1_J)J) \otimes S = (I \otimes S \cap S1_J)(J \otimes S) = I \otimes S$$

where the first equality is because S is faithfully flat and the second is by the previous case, since $I \otimes S$ is an outer ideal. It follows that $I = (I \cap R1_J)J$ as desired. \square

Remark. In the proof above, the inclusion $(I \cap R1_J)J \subseteq I$ could instead have been argued as follows. Define $\text{Sq}(J)$ as the R -submodule of J generated by x^2 for $x \in J$. Since 1_J is unimodular, one finds that $(I \cap R1_J)\text{Sq}(J) \subseteq I$. Then, one argues that $\text{Sq}(J) = J$ for a split Freudenthal algebra, and that $\text{Sq}(J \otimes S) = \text{Sq}(J) \otimes S$ for all flat $S \in R\text{-alg}$.

Corollary 8.4. *Every homomorphism $J \rightarrow J'$ of Freudenthal R -algebras is injective.*

Proof. Write ϕ for such a homomorphism. The kernel of ϕ is an ideal of J and therefore $\mathfrak{a}J$ for some ideal \mathfrak{a} of R . For $\alpha \in \mathfrak{a}$, we have $0 = \phi(\alpha 1_J) = \alpha \phi(1_J) = \alpha 1_{J'}$, so $\alpha = 0$ because $1_{J'}$ is unimodular (Lemma 6.12). $0 = \phi(\alpha 1_J) = \alpha \phi(1_J) = \alpha 1_{J'}$, so $\alpha = 0$ because $1_{J'}$ is unimodular. \square

Remark. There is also the notion of an *inner* ideal in a Jordan algebra, see [McC71b, Th. 8] for a description of them for $\text{Her}_3(\text{Zor}(R))$. The inner ideals are related to the projective homogeneous varieties associated with the group of isometries described in §14 and “outer automorphisms” relating these varieties, see [Rac77] and [CarrG].

9. GROUPS OF TYPE F_4 AND C_3

In the following, for a Jordan R -algebra J , we write $\text{Aut}(J)$ for the ordinary group of R -linear automorphisms of J and $\mathbf{Aut}(J)$ for the functor from $R\text{-alg}$ to groups such that $S \mapsto \text{Aut}(J \otimes S)$. Recall that for every simple root datum, there is a unique simple group scheme over \mathbb{Z} called a *Chevalley group* [DG70, Cor. XXIII.5.4], and every split simple

algebraic group over a field is obtained from a unique Chevalley group by base change [Mil17, §23g].

Lemma 9.1. *Let J be a Freudenthal algebra of rank 15 or 27 over a ring k . Then $\mathbf{Aut}(J)$ is a semisimple k -group scheme that is adjoint (i.e., its center is the trivial group scheme). Its root system has type C_3 if J has rank 15 and type F_4 if J has rank 27. If J is the split Freudenthal algebra, then the group scheme $\mathbf{Aut}(J)$ is obtained from the Chevalley group over \mathbb{Z} by base change.*

Proof. First suppose that $R = \mathbb{Z}$ and J is split. If J has rank 15, then the proof of 14.19 in [Spr73] shows that the automorphisms of $J \otimes F$ for every field F are exactly the automorphisms of the algebra $\mathrm{Mat}_6(F)$ with the split symplectic involution, which is the split adjoint group PGSp_6 . For J of rank 27, $\mathbf{Aut}(J) \times F$ is split of type F_4 by [Jac71, §6] (written for Lie algebras), [Fre85, Satz 4.11] (written for \mathbb{R}), [SV00, Th. 7.2.1] (if $\mathrm{char} F \neq 2, 3$), or [Spr73, 14.24] in general.

Note that $\mathbf{Aut}(J) \times F$ is connected and smooth as a group scheme over F , and $\mathbf{Aut}(J)$ is finitely presented (because \mathbb{Z} is noetherian and J is a finitely generated module), so it follows by [AlsG, Lemma B.1] that $\mathbf{Aut}(J)$ is smooth as a scheme over the Dedekind domain \mathbb{Z} . In summary, $\mathbf{Aut}(J)$ is semisimple and adjoint of the specified type. Moreover, because $\mathbf{Aut}(J) \times \mathbb{Q}$ is split, $\mathbf{Aut}(J)$ is a Chevalley group [Con, Th. 1.4].


In the case of general R and J , let $S \in R\text{-alg}$ be faithfully flat such that $J \otimes S$ is split. Then $\mathbf{Aut}(J) \times S$ is semisimple adjoint of the specified type. Certainly, $\mathbf{Aut}(J)$ is also smooth. Moreover, for each $\mathfrak{p} \in \mathrm{Spec} R$, there is a $\mathfrak{q} \in \mathrm{Spec} S$ such that $\mathfrak{q} \cap R = \mathfrak{p}$. Then the field of fractions $R(\mathfrak{p})$ of R/\mathfrak{p} embeds in the field $S(\mathfrak{q})$, so the algebraic closure $\overline{R(\mathfrak{p})}$ includes in the algebraic closure $\overline{S(\mathfrak{q})}$. Because $\mathbf{Aut}(J) \times \overline{S(\mathfrak{q})}$ is adjoint semisimple of the specified type and this property is unchanged by replacing one algebraically closed field by a smaller one, the same holds over $\overline{R(\mathfrak{p})}$. Since this holds for every \mathfrak{p} , the claim is verified. \square

Remark 9.2. In case R is a field, the automorphism group of the split Freudenthal algebra of rank 6 or 9 can be deduced in a similar manner, referring to 14.17 and 14.16 in [Spr73]. The automorphism group of the split Freudenthal algebra of rank 9 is $\mathrm{PGL}_3 \rtimes \mathbb{Z}/2$. The automorphism group of the split Freudenthal algebra of rank 6 is the special orthogonal group of the quadratic form $x^2 + y^2 + z^2$, i.e., the group commonly denoted $\mathrm{SO}(3)$. In particular, it is not smooth when R is a field of characteristic 2 and indeed one can give examples of Freudenthal algebras of rank 6 over a field of characteristic 2 that are not split by any étale cover.

For J, J_0 Jordan R -algebras, we define $\mathrm{Iso}(J, J_0)$ to be the set of R -linear isomorphisms $J \rightarrow J_0$ and $\mathbf{Iso}(J, J_0)$ to be the corresponding functor from $R\text{-alg}$ to sets defined by $S \mapsto \mathrm{Iso}(J \otimes S, J_0 \otimes S)$. If J and J_0 become isomorphic over a faithfully flat $S \in R\text{-alg}$, then $\mathbf{Iso}(J, J_0)$ is naturally an $\mathbf{Aut}(J_0)$ -torsor in the fpqc topology.

The statement of the following result is similar to statements over a field that can be found in [Ser02]. Its proof amounts to combining the lemma with the general machinery of descent.

Theorem 9.3. *Let J_0 be a Freudenthal R -algebra of rank $r = 15$ or 27 . In the diagram*



all arrows are bijections.

Proof. The facts that the arrows are well defined, the diagram commutes, and the diagonal arrows are injective are general feature of the machinery of descent. The lower left arrow is surjective because every Freudenthal algebra is split by some faithfully flat R -algebra by definition. The lower right arrow is surjective because every semisimple group scheme is split by some faithfully flat R -algebra (even an étale cover) [DG70, Cor. XXIV.4.1.6]. \square

In the theorem, the set $H^1(R, \mathbf{Aut}(J_0))$ is naturally a pointed set and the bijections are actually of pointed sets, where the distinguished elements are J_0 in the upper left and $\mathbf{Aut}(J_0)$ in the upper right.

In case R is a field of characteristic different from 2, 3 and $r = 27$, the theorem goes back to [Hij63]. Or see [KMRT98, 26.18].

Corollary 9.4. *For each Freudenthal R -algebra J of rank 15 or 27, there is an étale cover $S \in R\text{-alg}$ such that $J \otimes S$ is a split Freudenthal algebra.*

Proof. Let J_0 be the split Freudenthal R -algebra of the same rank as J . The image $\mathbf{Iso}(J, J_0)$ of J in $H^1(R, \mathbf{Aut}(J_0))$ is a $\mathbf{Aut}(J_0)$ -torsor. Since $\mathbf{Aut}(J_0)$ is smooth (Lemma 9.1), there is an étale cover of R that trivializes $\mathbf{Iso}(J, J_0)$. \square

Note that exactly the same kind of argument gives analogues of Lemma 9.1 and Theorem 9.3 for composition algebras, where $r = 4$ or 8 , and the group is of type A_1 or G_2 respectively.

10. GENERIC MINIMAL POLYNOMIAL OF A FREUDENTHAL ALGEBRA

Polynomials with polynomial-law coefficients. Let J be a Jordan R -algebra, $\mathcal{P}(J, R)$ the R -algebra of polynomial laws from J to R , and t a variable. Consider a polynomial $\mathbf{p}(t) = \sum_{i=0}^n f_i t^i$ with $f_i \in \mathcal{P}(J, R)$ for $0 \leq i \leq n$. For $S \in R\text{-alg}$, $x \in J \otimes S$, we have $\mathbf{p}(t, x) := \sum_{i=0}^n f_{iS}(x) t^i \in S[t]$, and we define

$$\mathbf{p}(x, x) := \sum_{i=0}^n f_{iS}(x) x^i \in J \otimes S.$$

The algebra J is said to *satisfy \mathbf{p}* if $\mathbf{p}(x, x) = 0 = (t\mathbf{p})(x, x)$ for all $x \in J \otimes S$, $S \in R\text{-alg}$. Note that the second equation follows from the first if 2 is invertible in R but not in general, see Remark 6.11.

The generic minimal polynomial. Let $J := \text{Her}_3(C, \Gamma)$ as in Example 6.13. With a variable t we recall from (6.10) that J satisfies the monic polynomial

$$(10.1) \quad \mathbf{m}_J = t^3 - \text{Tr}_J \cdot t^2 + S_J \cdot t - N_J \in \mathcal{P}(J)[t].$$

More precisely, by [Loo06, 2.4(b)], J is generically algebraic of degree 3 in the sense of [Loo06, 2.2] and \mathbf{m}_J is the *generic minimal polynomial* of J , i.e., the unique monic polynomial in $\mathcal{P}(J, R)[t]$ of minimal degree satisfied by J [Loo06, 2.7]. It follows that the Jordan algebra J determines the polynomial \mathbf{m}_J uniquely. In particular, the *generic norm* N_J , the *generic trace* T_J (or Tr_J) and, in fact, the cubic norm structure underlying J in the sense of Definition 6.1 are uniquely determined by J as a Jordan algebra.

By faithfully flat descent, every Freudenthal algebra J has a uniquely determined generic minimal polynomial of the form (10.1), and a uniquely determined underlying cubic norm structure. We conclude:

Proposition 10.2. *Every Freudenthal algebra is a cubic Jordan algebra.* □

The preceding discussion shows that, for a Freudenthal R -algebra J , the Jordan algebra structure of J alone (ignoring that J is a *cubic* Jordan algebra) determines the bilinear form T_J . (For example, the 11 Freudenthal \mathbb{R} -algebras discussed in Example 6.18 have distinct trace forms and therefore are distinct.) When R is a field of characteristic $\neq 2, 3$ and J and J' are reduced Freudenthal algebras, Springer proved that the converse also holds, i.e., $J \cong J'$ if and only if $T_J \cong T_{J'}$ [SV00, Th. 5.8.1]. We do not use Springer's result in this paper.

The following result can also be found in [Pet19, Cor. 18(b)], based on the different definition of Albert algebra appearing there.

Lemma 10.3. *Let J and J' be Freudenthal R -algebras. An R -linear map $\phi: J \rightarrow J'$ is an isomorphism of J and J' as Jordan algebras if and only if ϕ is surjective, $\phi(1_J) = 1_{J'}$, and $N_{J'} = N_J \phi$ as polynomial laws.*

Proof. The “only if” direction follows from the uniqueness of the generic minimal polynomial as in §10, so we show “if”. The equality $N_{J'} = N_J \phi$ of polynomial laws and the definition of the directional derivative in §3 gives formulas such as

$$\nabla_y N_J(x) = \nabla_{\phi(y)} N_{J'}(\phi(x)).$$

Since $\phi(1_J) = 1_{J'}$, the definition of the bilinear forms T_J and $T_{J'}$ in (6.2) give:

$$T_{J'}(\phi(x), \phi(y)) = T_J(x, y)$$

for all x, y . Therefore, on the one hand we have

$$\nabla_y N_J(x) = T_J(x^\sharp, y) = T_{J'}(\phi(x^\sharp), \phi(y)).$$

On the other hand, we have

$$\nabla_y N_J(x) = \nabla_{\phi(y)} N_{J'}(\phi(x)) = T_{J'}((\phi(x))^\sharp, \phi(y)).$$

Therefore, $\phi(x^\sharp) = \phi(x)^\sharp$ for all x . In summary, ϕ commutes with \sharp and preserves T_J . Therefore, by (6.6), ϕ is a homomorphism of Jordan algebras.

Suppose that x is in $\ker \phi$. Then for all $y \in J$, $T_J(x, y) = T_{J'}(\phi(x), \phi(y)) = 0$, so $x = 0$ since the bilinear form T_J is regular. Since ϕ is both surjective and injective, it is an isomorphism. □

11. BASIC CLASSIFICATION RESULTS FOR ALBERT ALGEBRAS

In the case where R is a field such as the real numbers, a finite field, a local field, or a global field, one can find in many places in the literature classifications of Albert algebras proved using techniques involving algebras as in [SV00, §5.8]. For such an R , groups of type F_4 can be classified using techniques from algebraic groups, such as in [PR94, Ch. 6] or [Gil19]. The two approaches are equivalent by Theorem 9.3.

Example 11.1 (Albert algebras over \mathbb{R}). Up to isomorphism, there are three Albert \mathbb{R} -algebras, namely the split one $\text{Her}_3(\text{Zor}(\mathbb{R}))$, $\text{Her}_3(\mathbb{O}, \langle 1, 1, -1 \rangle)$, and $\text{Her}_3(\mathbb{O})$. Rather than proving this in the language of Jordan algebras as in [AlbJ, Th. 10], one may leverage Theorem 9.3 as follows. The three algebras are pairwise non-isomorphic because their trace forms are (Example 6.18). At the same time, a computation in the Weyl group of F_4 as in [Ser02, §III.4.5], [BoroE, 14.1], or [AdT, Table 3] shows that $H^1(\mathbb{R}, \text{Aut}(\text{Her}_3(\text{Zor}(\mathbb{R}))))$ has three elements. That is, there are exactly three isomorphism classes of simple affine group schemes over \mathbb{R} of type F_4 , so we have found all of them.

Below, we will focus our attention on classification results in the case where R is not a field. We translate known results about cohomology of affine group schemes into the language of Albert algebras.

Proposition 11.2. *If R is (1) a complete discrete valuation ring whose residue field is finite or (2) a finite ring, then every Freudenthal R -algebra of rank 15 or 27 and every quaternion or octonion R -algebra is split.*

Proof. In view of Theorem 9.3 and its analogue for composition algebras, it suffices to prove that $H^1(R, \mathbf{G}) = 0$ for \mathbf{G} a simple R -group scheme of type F_4 or C_3 obtained by base change from a Chevalley group over \mathbb{Z} . In case (1), this is [Con, Prop. 3.10]. In case (2), we apply the following lemma. \square

Lemma 11.3. *If R is a finite ring and \mathbf{G} is a smooth connected R -group scheme, then $H^1(R, \mathbf{G}) = 0$.*

Proof. If R is not connected, then it is a finite product $R = \prod R_i$ where each ring R_i is finite, so $H^1(R, \mathbf{G}) = \prod H^1(R_i, \mathbf{G} \times R_i)$. Therefore it suffices to assume that R is connected.

Suppose \mathbf{X} is a \mathbf{G} -torsor. Our aim is to show that \mathbf{X} is the trivial torsor, i.e., $\mathbf{X}(R)$ is nonempty. Put \mathfrak{a} for the nil radical $\text{Nil}(R)$ of R . Because R is finite, there is some minimal $m \geq 1$ such that $\mathfrak{a}^m = 0$. We proceed by induction on m . If $m = 1$, then R is reduced and connected, so it is a finite field and $H^1(R, \mathbf{G}) = 0$ by Lang's Theorem. For the case $m \geq 2$, put $I := \mathfrak{a}^{m-1}$. The ring R/I has $\text{Nil}(R/I)^{m-1} = (\text{Nil}(R)/I)^{m-1} = 0$, so by induction $\mathbf{X}(R/I)$ is nonempty. On the other hand, $I^2 = \mathfrak{a}^{2m-2} = \mathfrak{a}^m \cdot \mathfrak{a}^{m-2} = 0$ and \mathbf{X} is smooth, so the natural map $\mathbf{X}(R) \rightarrow \mathbf{X}(R/I)$ is surjective. \square

Example 11.4. Suppose R is a Dedekind ring and write F for its field of fractions. For \mathbf{G} a Chevalley group of type G_2 , F_4 , or E_8 , the map $H^1(R, \mathbf{G}) \rightarrow H^1(F, \mathbf{G})$ has zero kernel [Har67, Satz 3.3]. Consequently, if A is an Albert or octonion R -algebra and $A \otimes F$ is split, then the R -algebra A is split.

In particular, if F is a global field with no real embeddings, then every Albert or octonion F -algebra is split, so every Albert or octonion R -algebra is split.

In the case where F is a number field with a real embedding, we provide the following partial result, which relies on Example 11.1.

Proposition 11.5. *Suppose F is a number field and R is a localization of its ring of integers at finitely many primes. If A is an Albert (resp., octonion) F -algebra such that $A \otimes \mathbb{R}$ is not isomorphic to $\text{Her}_3(\mathbb{O})$ (resp., \mathbb{O}) for every embedding $F \hookrightarrow \mathbb{R}$, then there is an Albert (resp., octonion) R -algebra B such that $B \otimes F \cong A$ and B is uniquely determined up to R -isomorphism.*

Proof. Write \mathbf{G} for the automorphism group of the split Albert (resp., octonion) F -algebra. Write $H_{\text{ind}}^1(R, \mathbf{G}) \subseteq H^1(R, \mathbf{G})$ for the isomorphism classes of R -algebras B such that $B \otimes F_v$ is not $\text{Her}_3(\mathbb{O})$ (resp., \mathbb{O}), i.e., such that $\mathbf{Aut}(B) \times F_v$ is not compact, for all real places v of F . Since \mathbf{G} is simply connected, Strong Approximation gives that the natural map $H_{\text{ind}}^1(R, \mathbf{G}) \rightarrow H_{\text{ind}}^1(F, \mathbf{G})$ is an isomorphism [Har67, Satz 4.2.4], which is what is claimed. \square

12. ISOTOPY

The aim of this section is to discuss the notion of isotopy of Jordan algebras, which will pay off later in the paper when we discuss groups of type E_6 in §14 and E_7 in §16. We include this material at this point in the paper because Corollary 12.9 is needed in the following section.

Definition 12.1. Let J be a Jordan R -algebra and suppose $u \in J$ is invertible. We define a Jordan algebra $J^{(u)}$ with the same underlying R -module, with U -operator $U_x^{(u)} = U_x U_u$ (where the unadorned U on the right denotes the U -operator in J), and with identity element $1^{(u)} = u^{-1}$. One checks that $J^{(u)}$ is indeed a Jordan algebra and for u, v invertible, we have $(J^{(u)})^{(v)} = J^{(U_u v)}$. A Jordan R -algebra J' is an *isotope* of J if it is isomorphic to $J^{(u)}$ for some invertible $u \in J$; equivalently one says that J and J' are *isotopic*. This defines an equivalence relation on Jordan algebras, which is a priori weaker than isomorphism.

We have presented the notion of isotopy here for Jordan algebras. However, there are analogous notions for other classes of algebras, which go back at least to [Alb 42]. For associative algebras, isotopy is the same as isomorphism. For octonion algebras, isotopy amounts to norm equivalence [AlsG, Cor. 6.7], which is a weaker condition than isomorphism, see [Gil14] and [AsHW].

Isotopes of cubic Jordan algebras. if J is a cubic Jordan R -algebra and $u \in J$ is invertible, then [McC69, Th. 2] and its proof show that the isotope $J^{(u)}$ is a cubic Jordan algebra as well whose identity element, adjoint and norm are given by

$$(12.2) \quad 1_{J^{(u)}} = u^{-1}, \quad x^{\sharp(u)} = N_J(u)U_u^{-1}x^{\sharp}, \quad N_{J^{(u)}}(x) = N_J(u)N_J(x).$$

Moreover, the (bi-)linear and quadratic trace of $J^{(u)}$ have the form

$$(12.3) \quad T_{J^{(u)}}(x, y) = T_J(U_u x, y), \quad \text{Tr}_{J^{(u)}}(x) = T_J(u, x), \quad S_{J^{(u)}}(x) = T_J(u^{\sharp}, x^{\sharp}).$$

The first equation of (12.3) is in [McC69, p. 500] while the second one follows from (12.2), the first, and Lemma 6.12 (1) via $\text{Tr}_{J^{(u)}}(x) = T_{J^{(u)}}(u^{-1}, x) = T_J(U_u u^{-1}, x) = T_J(u, x)$. Similarly,

$$S_{J^{(u)}}(x) = \text{Tr}_{J^{(u)}}(x^{\sharp(u)}) = T_J(u, N_J(u)U_u^{-1}x^{\sharp}) = T_J(N_J(u)U_u^{-1}u, x^{\sharp}) = T_J(u^{\sharp}, x^{\sharp}).$$

Example 12.4. $\text{Her}_3(C, \Gamma)$ is isotopic to $\text{Her}_3(C)$ for every Γ . Indeed, for

$$u := \begin{pmatrix} \gamma_1 & 0 & 0 \\ 0 & \gamma_2 & 0 \\ 0 & 0 & \gamma_3 \end{pmatrix} \in \text{Her}_3(C, \Gamma),$$

the map $\phi: \text{Her}_3(C, \Gamma)^{(u)} \rightarrow \text{Her}_3(C)$ defined by

$$\phi \begin{pmatrix} \alpha_1 & \gamma_2 c_3 & \gamma_3 \bar{c}_2 \\ \gamma_1 \bar{c}_3 & \alpha_2 & \gamma_3 c_1 \\ \gamma_1 c_2 & \gamma_2 \bar{c}_1 & \alpha_3 \end{pmatrix} = \begin{pmatrix} \gamma_1 \alpha_1 & \gamma_1 \gamma_2 c_3 & \cdot \\ \cdot & \gamma_2 \alpha_2 & \gamma_2 \gamma_3 c_1 \\ \gamma_1 \gamma_3 c_2 & \cdot & \gamma_3 \alpha_3 \end{pmatrix}$$

is an isomorphism of Jordan algebras. One can also turn this around:

$$\text{Her}_3(C, \Gamma) = (\text{Her}_3(C, \Gamma)^{(u)})^{(u^{-2})} \cong \text{Her}_3(C)^{(\phi(u^{-2}))} = \text{Her}_3(C)^{(u^{-1})}.$$

Jordan algebras isotopic to a split Freudenthal algebra. In the special case where R is a field, a Jordan algebra that is isotopic to the split Albert algebra $\text{Her}_3(\text{Zor}(R))$ is necessarily isomorphic to it, see for example [Jac71, p. 53, Th. 9]. Some hypothesis on R is necessary for the conclusion to hold. Alsaody has shown in [Als, Th. 2.7] that there exists a ring R finitely generated over \mathbb{C} and an Albert R -algebra that is isotopic to the split Albert R -algebra but is not isomorphic to it. We now show that it suffices to assume that R is local, see Corollary 12.8.

Theorem 12.5. *Suppose J is a Jordan R -algebra, where R is a local ring. If J is isotopic to $\text{Her}_3(C)$ for some composition R -algebra C , then J is isomorphic to $\text{Her}_3(C, \Gamma)$ for some Γ .*

Proof. By hypothesis, $J \cong \text{Her}_3(C)^{(u^{-1})}$ for some invertible $u \in \text{Her}_3(C)$. Example 12.4 shows we are done if u is diagonal.

Write N for the cubic form on $\text{Her}_3(C)$. In case u is not diagonal, we will apply successive elements $\eta \in \text{GL}(\text{Her}_3(C))$ such that $N\eta = N$ as polynomial laws. (In the notation of §14 below, $\eta \in \mathbf{Isom}(\text{Her}_3(C))(R)$.) Note that each such η defines an isomorphism of R -modules

$$(12.6) \quad \eta: \text{Her}_3(C)^{(u^{-1})} \rightarrow \text{Her}_3(C)^{(\eta(u)^{-1})}.$$

We have

$$N(\eta(u)^{-1}) = N(\eta(u))^{-1} = N(u)^{-1} = N(u^{-1}),$$

so we have by (12.2) that

$$N_{\text{Her}_3(C)^{(u^{-1})}} = N(u)^{-1}N = N(\eta(u)^{-1})N\eta = N_{\text{Her}_3(C)^{(\eta(u)^{-1})}}\eta.$$

Since η is a norm isometry that maps the identity element u^{-1} in the domain of (12.6) to the identity element in the codomain, it is an isomorphism of algebras by Lemma 10.3. Thus, if successive elements η transform u into a diagonal element, the proof will be complete.

We employ the transformation $\tau_{st}(q)$ for $1 \leq s \neq t \leq 3$ and $q \in C$ defined by

$$\tau_{st}(q) A \mapsto (I_3 + qE_{st})A(I_3 + \bar{q}E_{ts}),$$

where I_3 is the identity matrix, E_{st} is the 3-by-3 matrix with a 1 in the (s, t) -entry and 0 elsewhere, and juxtaposition defines naive multiplication of 3-by-3 matrices with entries in C . For example,

$$\tau_{12}(q) \begin{pmatrix} \alpha_1 & c_3 & \cdot \\ \cdot & \alpha_2 & c_1 \\ c_2 & \cdot & \alpha_3 \end{pmatrix} = \begin{pmatrix} \alpha_1 + \text{Tr}_C(q\bar{c}_3) + \alpha_2 N_C(q) & c_3 + \alpha_2 q & \cdot \\ \cdot & \alpha_2 & c_1 \\ c_2 + \bar{c}_1 \bar{q} & \cdot & \alpha_3 \end{pmatrix}.$$

These transformations appear in [Jac61, §5] and [Kru02, §2]; the argument in either reference shows that $\tau_{st}(q)$ preserves N for all choices of s, t , and q . Additionally, for every permutation π of $\{1, 2, 3\}$, there is a linear transformation that preserves N (actually, an automorphism of the algebra) that maps

$$(12.7) \quad \begin{pmatrix} \alpha_1 & c_3 & \cdot \\ \cdot & \alpha_2 & c_1 \\ c_2 & \cdot & \alpha_3 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_{\pi(1)} & c'_{\pi(3)} & \cdot \\ \cdot & \alpha_{\pi(2)} & c'_{\pi(1)} \\ c'_{\pi(2)} & \cdot & \alpha_{\pi(3)} \end{pmatrix},$$

where $c'_{\pi(i)}$ is a linear function of c_i for each i .

Write

$$u = \begin{pmatrix} \alpha_1 & c_3 & \cdot \\ \cdot & \alpha_2 & c_1 \\ c_2 & \cdot & \alpha_3 \end{pmatrix}.$$

By hypothesis $N(u)$ is invertible, i.e., does not lie in the maximal ideal \mathfrak{m} . We first argue that we may suppose $\alpha_1 \notin \mathfrak{m}$. If any α_i is invertible, then we may apply a transformation as in (12.7). If no α_i is invertible, then by (6.15) we have

$$N(u) \equiv \text{Tr}_C(c_1 c_2 c_3) \pmod{\mathfrak{m}},$$

whence $c_1 \notin \mathfrak{m}C$. Since n_C continues to be regular when changing scalars to R/\mathfrak{m} , some $q \in C$ has $n_C(q, c_1) \notin \mathfrak{m}$. Applying $\tau_{12}(q)$, we may arrange $\alpha_1 \notin \mathfrak{m}$.

Next we argue that we may assume that $c_2 = c_3 = 0$. We note that $\tau_{21}(q) \begin{pmatrix} \alpha_1 & c_3 & \cdot \\ \cdot & \alpha_2 & c_1 \\ c_2 & \cdot & \alpha_3 \end{pmatrix}$ has top row entries $\alpha_1, c_3 + \alpha_1 \bar{q}, \bar{c}_2$. Taking $q = -\bar{c}_3 a^{-1}$ shows that we may assume $c_2 = 0$. The argument that we may assume $c_3 = 0$ is similar, with the role of τ_{21} replaced by τ_{31} .

We have transformed u to an element of the form $\begin{pmatrix} \alpha_1 & 0 & \cdot \\ \cdot & \alpha_2 & c_1 \\ 0 & \cdot & \alpha_3 \end{pmatrix}$ of norm $\alpha_1(\alpha_2 \alpha_3 - N_C(c_1)) \notin \mathfrak{m}$, therefore at least one of α_2, α_3 , or $N_C(c_1)$ is not in \mathfrak{m} . The same argument as two paragraphs above, with $\tau_{i(i+1)}$ replaced by τ_{23} , shows that we may assume that $\alpha_2 \notin \mathfrak{m}$. The same argument as in the preceding paragraph, with τ_{21} replaced by τ_{32} , shows that we may assume that $c_1 = 0$. Thus, we have transformed u into a diagonal element, completing the proof. \square

Corollary 12.8. *Suppose J is a Jordan R -algebra over a local ring R . If J is isotopic to a split Freudenthal algebra whose rank does not take the value 6, then J is itself a split Freudenthal algebra.*

Proof. Combine the theorem and Proposition 7.4. \square

The hypothesis that J does not have rank 6 is necessary, because $\text{Her}_3(\mathbb{R}, \langle 1, 1, -1 \rangle)$ is isotopic to the split Freudenthal algebra $\text{Her}_3(\mathbb{R})$ (Example 12.4) but is not isomorphic to it (Example 6.18).

Corollary 12.9. *Every isotope of a Freudenthal algebra is itself a Freudenthal algebra.*

Proof. Suppose J is an isotope of a Freudenthal algebra. After base change to a faithfully flat extension, J is an isotope of a split Freudenthal algebra.

The R -algebra $S := \prod_{\mathfrak{m}} R_{\mathfrak{m}}$, where \mathfrak{m} ranges over maximal ideals of R , is faithfully flat. For each \mathfrak{m} , $J \otimes S_{\mathfrak{m}}$ is $\text{Her}_3(C, \Gamma)$ for C a split composition $S_{\mathfrak{m}}$ -algebra and some Γ by Theorem 12.5. By Proposition 7.2, there is a faithfully flat $S_{\mathfrak{m}}$ -algebra T such that $J \otimes T$ is a split Freudenthal algebra. The product of these T 's is a faithfully flat R -algebra over which J is the split Freudenthal algebra. \square

We close this section by making explicit the relationship between isotopy and norm similarity between Freudenthal algebras, extending Lemma 10.3.

Proposition 12.10. *Let J and J' be Freudenthal R -algebras. For an R -linear map $\phi : J \rightarrow J'$, the following are equivalent:*

- (1) ϕ is an isomorphism $J \rightarrow (J')^{(u)}$ for some invertible $u \in J'$ (“ ϕ is an isotopy”).
- (2) $N_{J'} \phi = \alpha N_J$ as polynomial laws for some $\alpha \in R^\times$, and ϕ is surjective (“ ϕ is a norm similarity”).

Proof. Since $(J')^{(u)}$ is a Freudenthal algebra by Corollary 12.9, condition (2) follows from (1) by Lemma 10.3 and (12.2). Conversely, we assume (2) and prove (1). Because $N_{J'}(\phi(1_J)) = \alpha$, the element $\phi(1_J)$ is invertible in J' . We set $u := \phi(1_J)^{-1}$ and $J'' := (J')^{(u)}$. We have

$$\phi(1_J) = u^{-1} = 1_{J''}.$$

Also, $N_{J'}(u) = N_{J'}(\phi(1_J))^{-1} = \alpha^{-1}$. Then

$$N_{J''}\phi = N_{J'}(u)N_{J'}\phi = N_J$$

as polynomial laws. Lemma 10.3 implies that ϕ is an isomorphism $J \xrightarrow{\sim} J''$, as desired. \square

13. CLASSIFICATION OF ALBERT ALGEBRAS OVER \mathbb{Z}

In this section, we study Albert algebras over the integers.

Definition 13.1. In the notation of Example 4.9, consider the element

$$\beta := (-1 + e_1 + e_2 + \cdots + e_7)/2 = h_1 + h_2 + h_3 - (2 + e_1) \in \mathcal{O},$$

as was done in [EG96, (5.2)]. That element has

$$\mathrm{Tr}_{\mathcal{O}}(\beta) = -1, \quad n_{\mathcal{O}}(\beta) = 2, \quad \text{and} \quad \beta^2 + \beta + 2 = 0.$$

Put

$$v := \begin{pmatrix} 2 & \beta & \cdot \\ \cdot & 2 & \beta \\ \beta & \cdot & 2 \end{pmatrix} \in \mathrm{Her}_3(\mathcal{O}).$$

Since $\mathrm{Tr}_{\mathcal{O}}(\beta^3) = 5$, we find that $N_{\mathrm{Her}_3(\mathcal{O})}(v) = 1$. In particular, v is invertible with inverse v^\sharp . We define $\Lambda := \mathrm{Her}_3(\mathcal{O})^{(v)}$; it is an Albert algebra by Corollary 12.9.

Proposition 13.2. $\mathrm{Her}_3(\mathcal{O}) \not\cong \Lambda$ as Jordan \mathbb{Z} -algebras, but $\mathrm{Her}_3(\mathcal{O}) \otimes \mathbb{Q} \cong \Lambda \otimes \mathbb{Q}$ as Jordan \mathbb{Q} -algebras.

Proof. We first prove the claim over \mathbb{Z} , which amounts to a computation from [EG96]. The isomorphism class of a Freudenthal algebra determines its cubic norm form and also its trace linear form. From (12.2) we deduce for $x \in \mathrm{Her}_3(\mathcal{O})$ that $x^\sharp(v) = 0$ if and only if $x^\sharp = 0$. Hence [EG96, Prop. 5.5] says that $\mathrm{Her}_3(\mathcal{O})$ contains exactly 3 elements x such that $x^\sharp = 0$ and $\mathrm{Tr}_{\mathrm{Her}_3(\mathcal{O})}(x) = 1$, whereas Λ has no elements x such that $x^\sharp(v) = 0$ and

$$T_{\mathrm{Her}_3(\mathcal{O})}(v, x) = 1,$$

where the left side is $\mathrm{Tr}_{\Lambda}(x)$ by (12.3). This proves that $\mathrm{Her}_3(\mathcal{O}) \not\cong \Lambda$.

Now consider $\mathrm{Her}_3(\mathcal{O}) \otimes \mathbb{R}$. It is called a ‘‘euclidean’’ Jordan algebra or, in older references, a ‘‘formally real’’ Jordan algebra, because every sum of nonzero squares is not zero [BrK, p. 331]. The element v has generic minimal polynomial, in the sense of (10.1), $(x-1)(x^2-5x+1)$, which has three positive real roots. Therefore, there is some $u \in \mathrm{Her}_3(\mathcal{O}) \otimes \mathbb{R}$ such that $u^2 = v$ [BrK, §XI.3, S. 3.6 and 3.7]. From this, it is trivial to see that

$$\Lambda \otimes \mathbb{R} \cong (\mathrm{Her}_3(\mathcal{O}) \otimes \mathbb{R})^{(v)} \cong \mathrm{Her}_3(\mathcal{O}) \otimes \mathbb{R}.$$

Since $\mathbf{G} := \mathbf{Aut}(\mathrm{Her}_3(\mathcal{O}))$ is simple and simply connected, the natural map $H^1(\mathbb{Q}, \mathbf{G}) \rightarrow H^1(\mathbb{R}, \mathbf{G})$ is a bijection, see [Har66] or [PR94, Th. 6.6]. Theorem 9.3 gives that $\mathrm{Her}_3(\mathcal{O}) \otimes \mathbb{Q} \cong \Lambda \otimes \mathbb{Q}$. \square

Theorem 13.3. *Over \mathbb{Z} :*

- (a) *There are exactly two isomorphism classes of octonion algebras: $\mathrm{Zor}(\mathbb{Z})$ and \mathcal{O} .*

- (b) *There are exactly four isomorphism classes of Albert algebras: $\text{Her}_3(\text{Zor}(\mathbb{Z}))$, $\text{Her}_3(\mathcal{O}, \langle 1, -1, 1 \rangle)$, $\text{Her}_3(\mathcal{O})$, and the algebra Λ .*
- (c) *There are exactly two isotopy classes of Albert algebras: $\text{Her}_3(\text{Zor}(\mathbb{Z}))$ and $\text{Her}_3(\mathcal{O})$.*

Proof. No pair of the listed algebras are isomorphic to another one. For $\text{Her}_3(\mathcal{O})$ and Λ , this is Prop. 13.2. For any other pair, base change to \mathbb{Q} yields non-isomorphic \mathbb{Q} -algebras.

Suppose that B is an octonion or Albert \mathbb{Z} -algebra. If B is indefinite, then it is determined by $B_{\mathbb{Q}}$ by Proposition 11.5. Since the indefinite octonion or Albert \mathbb{Q} -algebras are $\text{Zor}(\mathbb{Q})$, $\text{Her}_3(\text{Zor}(\mathbb{Q}))$, and $\text{Her}_3(\mathcal{O} \otimes \mathbb{Q}, \langle 1, -1, 1 \rangle)$, B is isomorphic to one of the algebras listed in the statement.

On the other hand, Gross’s mass formula [Gro96, Prop. 5.3] shows that there is only one composition \mathbb{Z} -algebra and two Albert \mathbb{Z} -algebras whose base change to \mathbb{Q} is definite. This shows that we have captured all the definite algebras as well, completing the proof of (a) and (b).

For (c), note that the three algebras in (b) that are not $\text{Her}_3(\text{Zor}(\mathbb{Z}))$ are all isotopic, see Example 12.4, so the two algebras listed in (c) represent all of the isotopy classes of Albert \mathbb{Z} -algebras. The base change of these two algebras to \mathbb{Q} have distinct co-ordinate algebras and therefore are not isotopic (Example 12.4), consequently they are not isotopic as \mathbb{Z} -algebras. \square

Note that part (a) of the theorem can be proved entirely in the language of octonion algebras, see [vdBS59].

In view of Theorem 9.3, part (b) is equivalent to a classification of the group schemes of type F_4 over \mathbb{Z} , which was done in Sections 6 and 7 of [Con], especially Examples 6.7 and 7.4. The innovation here is that we can use the language of Albert algebras also in the case of \mathbb{Z} where 2 is not invertible. Because of this extra flexibility, we can substitute results from the literature over algebraically closed fields (including characteristic 2) for some of the computations over \mathbb{Z} done in [Con].

Part (c) corresponds to the classification of groups of type E_6 over \mathbb{Z} up to isogeny, see §17.

Remark 13.4 (the Tits construction). The examples of Albert algebras exhibited so far have all been reduced algebras, i.e., Albert algebras of the kind described in Example 6.13. Such algebras are not division algebras, for example the element ε_i is not invertible. Historically speaking, it took many years after Albert algebras were defined — all the way until 1958 — for the first Albert division algebra to be exhibited in [Alb 58]. One reason for the difficulty is that, for there to exist an Albert division algebra over a field F of characteristic $\neq 3$, one needs $H^3(F, \mathbb{Z}/3) \neq 0$, see [Ros91], [PR96], or [Gar09, §8]. Conversely, if $H^3(F, \mathbb{Z}/3) \neq 0$, as happens when $F = \mathbb{Q}(t)$ for example, then one can construct an Albert division algebra via the so-called first Tits construction. This construction was first described in print in [Jac68, §IX.12] and later extended in various ways, including to the case of an arbitrary base ring in [PR86].

14. GROUPS OF TYPE E_6

Roundness of the norm. We note that the cubic norm of a Freudenthal algebra has the following special property. A quadratic form with this property is called “round”, see [EKM08, §9.A].

Lemma 14.1 (roundness). *For every Freudenthal R -algebra J ,*

$$\{\alpha \in R^\times \mid \alpha N_J \cong N_J\} = \{N_J(x) \in R^\times \mid x \text{ invertible in } J\}.$$

Proof. If $\alpha \in R^\times$ and $\phi \in \mathrm{GL}(J)$ are such that $\alpha N_J = N_J \phi$, then for $x := \phi(1_J)$ we have $N_J(x) = \alpha$. Conversely, if x is invertible in J , put $\alpha := N_J(x)$ and define $\phi := \alpha U_{x^{-1}}$. Then $N_J \phi = \alpha^3 N_J(x^{-1})^2 N_J$ by Lemma 6.12(3), so $N_J \phi = \alpha N_J$. \square

Example 14.2. For $J = \mathrm{Her}_3(C, \Gamma)$, the sets displayed in Lemma 14.1 equal R^\times . To see this for the right side, take $\alpha \in R^\times$ and note that $N_J(\alpha \varepsilon_1 + \varepsilon_2 + \varepsilon_3) = \alpha$. For the left side, consider $\phi \in \mathrm{GL}(J)$ defined by

$$\begin{aligned} \phi(\varepsilon_i) &= \alpha \varepsilon_i & \text{and} & & \phi(\delta_i(c)) &= \delta_i(c) & \text{for } i = 1, 2, \\ \phi(\varepsilon_3) &= \alpha^{-1} \varepsilon_3 & \text{and} & & \phi(\delta_3(c)) &= \delta_3(\alpha c). \end{aligned}$$

Then $N_J \phi = \alpha N_J$ as polynomial laws.

Example 14.3. In contrast to the preceding example, we now show that the sets displayed in Lemma 14.1 may be properly contained in R^\times . Suppose F is a field and J is a Freudenthal F -algebra such that N_J is anisotropic, i.e., $N_J(x) = 0$ if and only if $x = 0$. (For example, such a J exists if F is Laurent series or rational functions in one variable over a global field, see Remark 13.4.) We claim that, for t an indeterminate, every nonzero element in the image of $N_{J \otimes F[[t]]}$ has lowest term of degree divisible by 3. Because the norm is a homogeneous form, it suffices to prove this claim for $J \otimes F[[t]]$.

Let $x \in J \otimes F[[t]]$ be nonzero, so $x = \sum_{j \geq j_0} x_j t^j$ for some $j_0 \geq 0$ with $x_{j_0} \neq 0$. Since N_J is anisotropic, $N_J(x_0) \neq 0$. If $j_0 = 0$, then the homomorphism $F[[t]] \rightarrow F$ such that $t \mapsto 0$ sends $x \mapsto x_0$ and $N_{J \otimes F[[t]]}(x) \mapsto N_J(x_0) \neq 0$, therefore $N_{J \otimes F[[t]]}(x)$ has lowest degree term $N_J(x_0)t^0$. If $j_0 > 0$, then

$$N_{J \otimes F[[t]]}(x) = N_{J \otimes F[[t]]}(t^{j_0}(xt^{-j_0})) = t^{3j_0}(N_J(x_{j_0})t^0 + \dots),$$

proving the claim.

Corollary 14.4. *For Freudenthal R -algebras J and J' , the following are equivalent:*

- (1) J and J' are isotopic.
- (2) $N_J \cong \alpha N_{J'}$ for some $\alpha \in R^\times$.
- (3) $N_J \cong N_{J'}$.

Proof. The equivalence of (1) and (2) is Proposition 12.10.

Supposing (2), let $\phi: J' \rightarrow J$ be an R -module isomorphism such that $\alpha N_{J'} = N_J \phi$. Take $x := \phi(1_{J'})$. Since $N_J(x) = \alpha$, Lemma 14.1 gives that $\alpha N_{J'} \cong N_J$. As N_J is also isomorphic to $\alpha N_{J'}$, we conclude (3). The converse is trivial. \square

In the corollary, the inclusion of (3) seems to be new, even in the case where R is a field. Omitting that, in the special case where R is a field of characteristic $\neq 2, 3$, the equivalence of (1) and (2) and Proposition 14.7 below can be found as Theorems 7 and 10 in [Jac71].

Albert algebras and groups of type E_6 . The stabilizer of the cubic form N_J in $\mathbf{GL}(J)$ is a closed sub-group-scheme denoted $\mathbf{Isom}(J)$. It contains $\mathbf{Aut}(J)$ as a natural sub-group-scheme. Arguing as in the proof of Lemma 9.1, one finds that $\mathbf{Isom}(J)$ is a simple affine group scheme that is simply connected of type E_6 . (In the case where R is an algebraically closed field, this claim is verified in [Spr73, 11.20, 12.4], or see [SV00, Th. 7.3.2] for the case where R is a field of characteristic different from 2, 3.) Compare [Als, Lemma 2.3] or [Con, App. C]. Moreover, $\mathbf{Isom}(J)$ is a ‘‘pure inner form’’ in the sense of [Con, §3], resp. ‘‘strongly inner’’ in [CalF, Def. 2.2.4.9], meaning that it is obtained by twisting the group scheme $\mathbf{Isom}(J_0)$ for the split Albert algebra J_0 by a class with values in $\mathbf{Isom}(J)$.

We note that the center of $\mathbf{Isom}(J)$ is the group scheme μ_3 of cube roots of unity operating on J by scalar multiplication

Faithfully flat descent shows that the set $H^1(R, \mathbf{Isom}(J))$ is in bijection with isomorphism classes of pairs (M, f) , where M is a projective module of the same rank as J and f is a cubic form on M — i.e., an element of $\mathbf{S}^3(M^*)$ — such that $f \otimes S$ is isomorphic to the norm on $\mathrm{Her}_3(\mathrm{Zor}(S))$ for some faithfully flat $S \in R\text{-alg}$. For every Albert R -algebra J and every $\alpha \in R^\times$, $(J, \alpha N_J)$ is such a pair by Example 14.2. In the special case where R is a field, every such pair (M, f) — i.e., every element of $H^1(R, \mathbf{Isom}(J))$ — is of the form $(J, \alpha N_J)$ for some J and $\alpha \in R^\times$, see [Gar09, 9.12] in general or [Spr62] for the case of characteristic $\neq 2, 3$.

Outer automorphism of $\mathbf{Isom}(J)$. Suppose J and J' are Freudenthal R -algebras and $\phi: J \rightarrow J'$ is an isomorphism of R -modules. Since the bilinear form $T_{J'}$ is regular, there is a unique R -linear map $\phi^\dagger: J \rightarrow J'$ such that $T_{J'}(\phi x, \phi^\dagger y) = T_J(x, y)$ for all $x, y \in J$. Because T_J and $T_{J'}$ are symmetric, we have $(\phi^\dagger)^\dagger = \phi$ for all ϕ . If J'' is another Freudenthal R -algebra and $\psi: J' \rightarrow J''$ is an R -linear bijection, then $(\phi\psi)^\dagger = \phi^\dagger\psi^\dagger$.

Proposition 14.5. *Let J be a Freudenthal R -algebra.*

- (1) *If $\phi \in \mathrm{GL}(J)$ is such that $N_J\phi = \alpha N_J$ for some $\alpha \in R^\times$, then $N_J\phi^\dagger = \alpha^{-1}N_J$.*
- (2) *The map $\phi \mapsto \phi^\dagger$ is an automorphism of $\mathbf{Isom}(J)$ of order 2 that is not an inner automorphism.*
- (3) *For ϕ as in (1) or in $\mathbf{Isom}(J)$, $\phi^\dagger = \phi$ if and only if ϕ is an automorphism of J .*

Proof. (1): Put $u := \phi(1_J)^{-1}$. On the one hand,

$$T_J(x, y) = T_{J^{(u)}}(\phi(x), \phi(y))$$

for all $x, y \in J$, because ϕ is an isomorphism $J \rightarrow J^{(u)}$ by Proposition 12.10. On the other hand, (12.3) yields

$$T_{J^{(u)}}(\phi(x), \phi(y)) = T_J(U_u\phi(x), \phi(y)).$$

Therefore,

$$(14.6) \quad \phi^\dagger = U_{\phi(1_J)}^{-1}\phi.$$

To complete the proof of (1), we note by Lemma 6.12(3) that

$$N_J\phi^\dagger = N_JU_u\phi = N_J(u)^2N_J\phi = \alpha^{-1}N_J.$$

For (2), we only have to check that the map is not an inner automorphism. Let $S \in R\text{-alg}$ be such that there exists $\zeta \in \mu_3(S)$ such that $\zeta \neq 1$. Then $\zeta^\dagger = \zeta^{-1} \neq \zeta$ and ζ is in the center of $\mathbf{Iso}(J)$, proving that the automorphism is not inner (and not the identity).

For (3), suppose $\phi^\dagger = \phi$. Then $N_J\phi = N_J$. By (14.6), $U_{\phi(1_J)} = \mathrm{Id}_J$, so $\phi(1_J) = \zeta 1_J$ for some $\zeta \in R$ with $\zeta^2 = 1$ (Example 7.3). Yet $1 = N_J(1_J) = N_J\phi(1_J)$, so ζ^3 also equals 1, whence $\phi(1_J) = 1_J$. Lemma 10.3 shows that ϕ is an automorphism of J . Conversely, if ϕ is an automorphism of J , then $u = 1_J$, so $\phi^\dagger = \phi$ by (14.6). \square

Proposition 14.7. *Let J and J' be Albert R -algebras. Among the statements*

- (1) $\mathbf{Isom}(J) \cong \mathbf{Isom}(J')$.
- (2) *There is a line bundle L and isomorphism $h: L^{\otimes 3} \rightarrow R$ such that $(J', N_{J'}) \cong [L, h] \cdot (J, N_J)$ for \cdot as defined in §3.*
- (3) *J and J' are isotopic.*

we have the implications (1) \Leftrightarrow (2) \Leftarrow (3). If $\mathrm{Pic} R$ has no 3-torsion other than zero, then all three statements are equivalent.

Proof. Suppose (1); we prove (2). We may assume R is connected.

The conjugation action gives a homomorphism $\mathbf{Isom}(J) \rightarrow \mathbf{Aut}(\mathbf{Isom}(J))$, which gives a map of pointed sets

$$(14.8) \quad H^1(R, \mathbf{Isom}(J)) \rightarrow H^1(R, \mathbf{Aut}(\mathbf{Isom}(J))),$$

where the second set is in bijection with isomorphism classes of R -group schemes that become isomorphic to $\mathbf{Isom}(J)$ after base change to an fppf R -algebra. By hypothesis, the class of $N_{J'} \in H^1(R, \mathbf{Isom}(J))$ is in the kernel of (14.8).

There is an exact sequence

$$1 \rightarrow \mathbf{Isom}(J)/\mu_3 \rightarrow \mathbf{Aut}(\mathbf{Isom}(J)) \rightarrow \mathbb{Z}/2 \rightarrow 1$$

of fppf sheaves by [DG70, Th. XXIV.1.3]. Since R is connected, $(\mathbb{Z}/2)(R)$ has one non-identity element, and it is the image of the map \dagger from Lemma 14.5. That is, in the exact sequence

$$\mathbf{Aut}(\mathbf{Isom}(J))(R) \rightarrow (\mathbb{Z}/2)(R) \rightarrow H^1(R, \mathbf{Isom}(J)/\mu_3) \rightarrow H^1(R, \mathbf{Aut}(\mathbf{Isom}(J))),$$

the first map is surjective, so the third map has zero kernel and we deduce that the image of $N_{J'}$ in $H^1(R, \mathbf{Isom}(J)/\mu_3)$ is the zero class. It follows that $N_{J'}$ is in the image of the map

$$H^1(R, \mu_3) \rightarrow H^1(R, \mathbf{Isom}(J)),$$

which is the orbit of the zero class N_J under the action of the group $H^1(R, \mu_3)$, which is (2).

That (2) implies (1) is Lemma 3.6. The claimed implications between (3) and (2) are Corollary 14.4. \square

15. FREUDENTHAL TRIPLE SYSTEMS

In this section, we define Freudenthal triple systems, also known as FT systems. We will see in Theorem 16.4 in the next section that they play the same role relative to groups of type E_7 that forms of the norm on an Albert algebra play for groups of type E_6 .

For any Albert R -algebra J , define $Q(J)$ to be the rank 56 projective R -module $R \oplus R \oplus J \oplus J$ endowed with a 4-linear form Ψ and an alternating bilinear form b , defined as follows.

We write a generic element of $Q(J)$ as $\begin{pmatrix} \alpha & x \\ x' & \alpha' \end{pmatrix}$ for $\alpha, \alpha' \in R$ and $x, x' \in J$. We define

$$(15.1) \quad b_J \left(\begin{pmatrix} \alpha & x \\ x' & \alpha' \end{pmatrix}, \begin{pmatrix} \beta & y \\ y' & \beta' \end{pmatrix} \right) := \alpha\beta' - \alpha'\beta + T_J(x, y') - T_J(x', y).$$

As an intermediate step to defining Ψ , define a quartic form

$$(15.2) \quad q_J \left(\begin{pmatrix} \alpha & x \\ x' & \alpha' \end{pmatrix} \right) = -4T_J(x^\sharp, x'^\sharp) + 4\alpha N_J(x) + 4\alpha' N_J(x') + (T_J(x, x') - \alpha\alpha')^2,$$

compare [Brown, p. 87] or [Kru07, p. 940].

To define the 4-linear form, consider first the case $R = \mathbb{Z}$ and $J := \text{Her}_3(\text{Zor}(\mathbb{Z}))$. (The following definitions are inspired by [Lur01, §6].) Putting X_i for an element of $Q(J)$ and t_i for an indeterminate, the coefficient of $t_1 t_2 t_3 t_4$ in $q(\sum t_i X_i)$, equivalently, the 4-linear form

$$(X_1, X_2, X_3, X_4) \mapsto \nabla_{X_1} \nabla_{X_2} \nabla_{X_3} q(X_4)$$

on $Q(J)$, equals 2Θ for a symmetric 4-linear form Θ . Define 4-linear forms Φ_i via

$$(15.3) \quad \Phi_1(X_1, X_2, X_3, X_4) = b(X_1, X_2) b(X_3, X_4)$$

$$(15.4) \quad \Phi_2(X_1, X_2, X_3, X_4) = b(X_1, X_3) b(X_4, X_2)$$

$$(15.5) \quad \Phi_3(X_1, X_2, X_3, X_4) = b(X_1, X_4) b(X_2, X_3).$$

Then $\Theta + \sum \Phi_i$ is divisible by 2 as a 4-linear function on $Q(\text{Zor}(\mathbb{Z}))$ and we set

$$(15.6) \quad \Psi_{\text{Her}_3(\text{Zor}(\mathbb{Z}))} := \frac{1}{2}(\Theta + \sum \Phi_i).$$

As Θ is symmetric, Ψ is evidently stable under even permutations of its arguments, and we have:

$$\Psi(X_1, X_2, X_3, X_4) - \Psi(X_2, X_1, X_3, X_4) = \sum \Phi_i.$$

For any ring R , we define $\Psi_{\text{Her}_3(\text{Zor}(R))} := \Psi_{\text{Her}_3(\text{Zor}(\mathbb{Z}))} \otimes R$, and we define Ψ_J for an Albert R -algebra J by descent.

Definition 15.7. A *Freudenthal triple system*¹ or *FT system* (M, Ψ, b) is an R -module M endowed with a 4-linear form Ψ and an alternating bilinear form b , such that $(M, \Psi, b) \otimes S$ is isomorphic (in an obvious sense) to $Q(J)$ for some faithfully flat $S \in R\text{-alg}$ and some Albert S -algebra J .

Comparison with other definitions. Suppose for this paragraph that 6 is invertible in R . Given an FT system (M, Ψ, b) , we may define 4-linear forms Φ_i on M via (15.4) and recover Θ and q via

$$(15.8) \quad \Theta := 2\Psi - \sum \Phi_i \quad \text{and} \quad \Theta(X, X, X, X) = 12q(X)$$

as polynomial laws in X . (This last is a special case of the general fact that going from a homogeneous form of degree d to a d -linear form and back to a homogeneous form of degree d equals multiplication by $d!$ [Bour, §IV.5.8, Prop. 12(i)].) Since the form b is regular and Θ is symmetric, the equation

$$\Theta(X_1, X_2, X_3, X_4) = b(X_1, t(X_2, X_3, X_4))$$

implicitly defines a symmetric 3-linear form $t: M \times M \times M \rightarrow M$, and $\mathbf{Aut}(M, \Psi, b)$ equals $\mathbf{Aut}(M, t, b)$. That is, under the hypothesis that 6 is invertible in R , we would obtain an equivalent class of objects if we replaced the asymmetric 4-linear form Ψ in the definition of FT systems with the quartic form q (the version studied in [Brown]) or with the trilinear form t (the version studied in [Mey68]).

Similarity of FT systems. For a d -linear form f on an R -module M , i.e., an R -linear map $f: M^{\otimes d} \rightarrow R$, and a d -trivialized line bundle $[L, h] \in H^1(R, \mu_d)$, we define a d -linear form $[L, h] \cdot f$ on $M \otimes L$ via the composition

$$(M \otimes L)^{\otimes d} \xrightarrow{\sim} M^{\otimes d} \otimes L^{\otimes d} \xrightarrow{f \otimes h} R.$$

For $Q := (M, \Psi, b)$ an FT system and a discriminant module $[L, h] \in H^1(R, \mu_2)$, we define $[L, h] \cdot Q$ to be the triple consisting of the module $M \otimes L$, the 4-linear form $[L, h^{\otimes 2}] \cdot \Psi$ for $[L, h^{\otimes 2}] \in H^1(R, \mu_4)$, and the bilinear form $[L, h] \cdot b$. Since $\langle 1 \rangle \cdot Q$ is Q itself, we deduce that $[L, h] \cdot Q$ is also an FT system. We say that FT systems Q, Q' are *similar* if $Q' \cong [L, h] \cdot Q$ for some $[L, h] \in H^1(R, \mu_2)$. For example, for any FT system (M, Ψ, b) and any $\alpha \in R^\times$, (M, Ψ, b) and $(M, \alpha^2 \Psi, \alpha b)$ are similar.

Example 15.9. Suppose $(M, \Psi, b) = Q(J)$ for some Albert R -algebra J . Then for every $\mu \in R^\times$, the map

$$\begin{pmatrix} \alpha & x \\ x' & \alpha' \end{pmatrix} \mapsto \begin{pmatrix} \alpha/\mu & \mu x \\ x' & \mu^2 \alpha' \end{pmatrix}$$

is an isomorphism $\langle \mu \rangle \cdot Q(J) \xrightarrow{\sim} Q(J)$. One checks this for $R = \mathbb{Z}$ and $J = \text{Her}_3(\text{Zor}(\mathbb{Z}))$ using (15.1) and (15.2). It follows for general R and J by base change and twisting.

¹See p. 273 of [Spr06] for remarks on the history of this term.

16. GROUPS OF TYPE E_7

We will now relate FT systems as defined in the previous section to affine group schemes of type E_7 . Here is a tool that allows us to work with the quartic form q as in (15.2) rather than the less-convenient 4-linear form Ψ , while still getting results that hold when 6 is not invertible.

Lemma 16.1. *Let (M, Ψ, b) be an FT system over \mathbb{Z} , let \mathbf{G} be a closed subgroup of $\mathbf{GL}(M)$, and let F be a field of characteristic zero. If $\mathbf{G}(F)$ is dense in \mathbf{G} (which holds if \mathbf{G} is connected) and $\mathbf{G}(F)$ preserves $b \otimes F$ and the quartic form q over F defined by (15.8), then G is a closed sub-group-scheme of $\mathbf{Aut}(M, \Psi, b)$.*

Proof. Since $\mathbf{G}(F)$ is dense in \mathbf{G} , the group scheme $\mathbf{G} \times F$ preserves $b \otimes F$ and q , whence also $\Psi \otimes F$. Viewing b and Ψ as elements of the representation $V := (M^*)^{\otimes d}$ of \mathbf{G} for $d = 2$ or 4 , the natural map $V^{\mathbf{G}} \otimes F \rightarrow (V \otimes F)^{\mathbf{G} \times F}$ is an isomorphism because F is flat over \mathbb{Z} [Ses77, Lemma 2], so \mathbf{G} preserves b and Ψ . \square

Corollary 16.2. *For every Freudenthal R -algebra J , there is an inclusion $f: \mathbf{Isom}(J) \hookrightarrow \mathbf{Aut}(Q(J))$ via*

$$f(\phi) \begin{pmatrix} \alpha & x \\ x' & \alpha' \end{pmatrix} = \begin{pmatrix} \alpha & \phi(x) \\ \phi^\dagger(x') & \alpha' \end{pmatrix}.$$

Proof. Consider the case $J = \text{Her}_3(\text{Zor}(\mathbb{Z}))$. For $\phi \in \mathbf{Isom}(J)(\mathbb{Q})$, it follows from the definition of ϕ^\dagger and Proposition 14.5(1) that $f(\phi)$ is an isomorphism of the bilinear and quartic forms $b \otimes \mathbb{Q}$ and q defined by (15.2) for $J \otimes \mathbb{Q}$. The lemma gives the claim in this case. Base change and twisting gives the claim for every R and every Albert R -algebra J . \square

Corollary 16.3. *Suppose J and J' are Albert R -algebras. If J and J' are isotopic, then $\mathbf{Aut}(Q(J)) \cong \mathbf{Aut}(Q(J'))$.*

Proof. The inclusions $\mathbf{Aut}(J) \hookrightarrow \mathbf{Isom}(J) \hookrightarrow \mathbf{Aut}(Q(J))$ induce maps

$$H^1(R, \mathbf{Aut}(J)) \rightarrow H^1(R, \mathbf{Isom}(J)) \rightarrow H^1(R, \mathbf{Aut}(Q(J))),$$

where the last set classifies FT systems over R . The class of J' in $H^1(R, \mathbf{Aut}(J))$ maps to the class of $N_{J'}$ in $H^1(R, \mathbf{Isom}(J))$, and by hypothesis and by Proposition 14.7 this is the trivial class. Therefore, the image of J' in $H^1(R, \mathbf{Aut}(Q(J)))$, which is $Q(J')$, is the trivial class. \square

In case R is a field of characteristic $\neq 2, 3$, the converse of Corollary 16.3 is true by [Fer72, Cor. 6.9]. That is, if $Q(J) \cong Q(J')$, then J and J' are isotopic.

Theorem 16.4. *The group scheme $\mathbf{Aut}(Q(\text{Her}_3(\text{Zor}(R))))$ over R is obtained from the simply connected Chevalley group of type E_7 over \mathbb{Z} by base change. Every strongly inner and simply connected simple R -group scheme of type E_7 over R is of the form $\mathbf{Aut}(Q)$ for some FT system Q . For FT systems Q and Q' , $\mathbf{Aut}(Q) \cong \mathbf{Aut}(Q')$ if and only if Q and Q' are similar.*

Proof. Put $J_R := \text{Her}_3(\text{Zor}(R))$ and $Q_R := Q(J_R)$. We will show that $\mathbf{Aut}(Q_R)$ is isomorphic to the base change to R of the simply connected Chevalley group E_7 over \mathbb{Z} .

In addition to the sub-group-scheme $\mathbf{Isom}(J_R)$ of $\mathbf{Aut}(Q_R)$ provided by Corollary 16.2, we consider \mathbb{G}_m defined by

$$\beta \begin{pmatrix} \alpha & x \\ x' & \alpha' \end{pmatrix} = \begin{pmatrix} \beta^{-3}\alpha & \beta x \\ \beta^{-1}x' & \beta^3\alpha' \end{pmatrix} \quad \text{for } \beta \in R^\times$$

and two copies of J_R (as group schemes under addition) through which an element $y \in J_R$ acts via

$$y \begin{pmatrix} \alpha & x \\ x' & \alpha' \end{pmatrix} = \begin{pmatrix} \alpha+b(x',y) & x+\alpha'y \\ x'+x \times y & \alpha' \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} \alpha & x+x' \times y \\ x'+\alpha y & \alpha'+b(x',y) \end{pmatrix}.$$

These preserve b and q , see for example [Brown, p. 95] or [Kru07, p. 942], and so by Lemma 16.1 do belong to $\mathbf{Aut}(Q_R)$. Considering the Lie algebras of $\mathbf{Isom}(J_R)$, \mathbb{G}_m , and the two copies of J , as subalgebras of $\mathrm{Lie}(\mathbf{GL}(Q_R))$, one can identify the subalgebra L_R they generate with the Lie algebra of $E_7 \times R$ by picking out specific root subalgebras and so on as in [Fre54] or [Sel63], or see [Gar01, §7] for partial information. Note that $\mathrm{Lie}(\mathbf{Aut}(Q_R)) \supseteq L_R$. For F any algebraically closed field, we may identify the smooth closed subgroup of $\mathbf{Aut}(Q_F)$ generated by $\mathbf{Isom}(J_F)$, \mathbb{G}_m , and the two copies of J_F with $E_7 \times F$.

In [Lur01], Lurie begins with $L_{\mathbb{Z}}$ and defines $L_{\mathbb{Z}}$ -invariant 4-linear forms Θ^L , Φ_i^L , and Ψ^L and alternating bilinear form b^L on the 56-dimensional Weyl module of $L_{\mathbb{Z}}$. Over \mathbb{C} , $\mathbf{Aut}(Q_{\mathbb{C}})$ is simply connected of type E_7 by the references in the previous paragraph, so it preserves the base change of Lurie's forms $\Theta^L \otimes \mathbb{C}$, etc. Because $\mathbf{Aut}(Q_{\mathbb{C}})(\mathbb{C})$ is dense in $\mathbf{Aut}(Q_{\mathbb{C}})$, Lemma 16.1 shows that $\mathbf{Aut}(Q_{\mathbb{Z}})$ preserves Θ^L , the Φ_i^L , Ψ^L , and b^L . By the uniqueness of E_7 invariant bilinear and symmetric 4-linear forms on M (which follows from the uniqueness over \mathbb{C} as in the proof of Lemma 16.1), we find that $b^L = \pm b$ and $\Theta^L = \pm \Theta$. Note that regardless of the sign on b in the preceding sentence, we find $\Phi_i^L = \Phi_i$ for all i and $\mathbf{Aut}(Q_F)$ preserves b^L . Now let F be an algebraically closed field. If F has characteristic different from 2, then $\mathbf{Aut}(Q_F)$ preserves $2\Psi = \Theta + \sum \Phi_i$ and the Φ_i , so it preserves Θ , hence Θ^L , hence Ψ^L . If F has characteristic 2, then although $\Psi^L = \frac{1}{2}(\pm\Theta + \sum \Phi_i)$ for some choice of sign as polynomials over \mathbb{Z} , we have $\Psi^L \otimes F = \Psi \otimes F$. In either case, $\mathbf{Aut}(Q_F)$ preserves $b^L \otimes F$ and $\Psi^L \otimes F$, whence so does its Lie algebra, so $\dim \mathrm{Lie} \mathbf{Aut}(Q_F) \leq \dim L_F$ by [Lur01, Th. 6.2.3]. Putting this together with the previous paragraph, we see that $\mathbf{Aut}(Q_F)$, an affine group scheme over the field F , is smooth with identity component $E_7 \times F$.

We claim that $\mathbf{Aut}(Q_F)$ is connected. Since its identity component E_7 has no outer automorphisms, every element of $\mathbf{Aut}(Q_F)(F)$ is a product of an element of $E_7(F)$ and a linear transformation centralizing E_7 . The action of $E_7 \times F$ on Q_F is irreducible (it is the 56-dimensional minuscule representation), so the centralizer of E_7 consists of scalar transformations. Finally, we note that the intersection of $\mathbf{Aut}(Q_F)$ and the scalar transformations is the group scheme μ_2 of square roots of unity, which is contained in E_7 . In summary, $\mathbf{Aut}(Q_F) = E_7 \times F$ for every algebraically closed field F .

As in the proof of Lemma 9.1, it follows that $\mathbf{Aut}(Q_{\mathbb{Z}})$ is a simple affine group scheme that is simply connected of type E_7 , and we deduce from the fact that $\mathbf{Aut}(Q_{\mathbb{R}})$ is split that $\mathbf{Aut}(Q_{\mathbb{Z}})$ is in fact the Chevalley group.

The second claim now follows by descent.

The third claim is proved in the same manner as Proposition 14.7, although the current situation is somewhat easier due to the absence of nontrivial automorphisms of the Dynkin diagram of E_7 and therefore the absence of outer automorphisms for semisimple groups of that type. Therefore, the sequence

$$(16.5) \quad H^1(R, \mu_2) \rightarrow H^1(R, \mathbf{Aut}(Q)) \rightarrow H^1(R, \mathbf{Aut}(\mathbf{Aut}(Q)))$$

is exact, where μ_2 is the center of $\mathbf{Aut}(Q)$ and $\mathbf{Aut}(\mathbf{Aut}(Q)) \cong \mathbf{Aut}(Q_R)/\mu_2$ is the adjoint group. We have $\mathbf{Aut}(Q') \cong \mathbf{Aut}(Q)$ if and only if the element Q' in $H^1(R, \mathbf{Aut}(Q))$ is in the kernel of the second map in (16.5), if and only if Q' is in the image of the first

map. To complete the proof, it suffices to calculate by descent that the action of $H^1(R, \mu_2)$ on $H^1(R, \mathbf{Aut}(Q))$ is exactly by the similarity action defined in §15. \square

Corollary 16.6. *If R is (1) a complete discrete valuation ring whose residue field is finite; (2) a finite ring; or (3) a Dedekind domain whose field of fractions F is a global field with no real embeddings, then the the split FT system is the only one over R , up to isomorphism.*

Proof. Imitate the arguments in Proposition 11.2 or Example 11.4, where \mathbf{G} is the base change to R of the simply connected Chevalley group $\mathbf{Aut}(Q(\mathrm{Zor}(\mathbb{Z})))$. \square

Remarks. A previous work that considered groups of type E_7 over rings is [Luz13]. Aschbacher [Asch] studied the 4-linear form in the case where R is a field of characteristic 2. The paper [MW19] studied the case of fields of any characteristic, organized around a polynomial law $\Theta \in \mathcal{P}(Q, R)$ that is not homogeneous.

For a field F of characteristic $\neq 2, 3$, FT systems have been studied in this century in [CI], [Hel12], [Kru07], [Spr06], and [BorsDF⁺] to name a few. They arise naturally in the context of the bottom row of the magic triangle from [DG02, Table 2], in connection with the existence of extraspecial parabolic subgroups as in [Röh93] or [Gar09, §12], or from groups with a BC_1 grading [GG21, p. 995]. For every Albert F -algebra J , the group scheme $\mathbf{Aut}(Q(J))$ is isotropic, see for example [Spr06, Lemma 5.6(i)]. Yet there exist strongly inner groups of type E_7 that are anisotropic, see [Tit90, 3.1] or [Gar09, App. A], and therefore there exist FT systems Q that are not isomorphic to $Q(J)$ for any J . A construction that produces all FT systems can be obtained by considering a subgroup $\mathrm{Isom}(J) \rtimes \mu_4$ of $\mathbf{Aut}(Q(J))$, which leads to a surjection $H^1(F, \mathrm{Isom}(J) \rtimes \mu_4) \rightarrow H^1(F, \mathbf{Aut}(Q(J)))$, see [Gar09, 12.13], [Gar01, Lemma 4.15], or [Spr06, §4].

17. EXCEPTIONAL GROUPS OVER \mathbb{Z}

We now give an explicit description of the isomorphism classes of semisimple affine group schemes over \mathbb{Z} of types F_4 , G_2 , E_6 , and E_7 .

There are four such group schemes of type F_4 , namely $\mathbf{Aut}(J)$ for each of the four Albert \mathbb{Z} -algebras listed in Theorem 13.3(b). The proof of this fact is intertwined with the proof of that theorem. Similarly, there are two such group schemes of type G_2 , namely $\mathbf{Aut}(C)$ for $C = \mathrm{Zor}(\mathbb{Z})$ or \mathcal{O} .

Proposition 17.1. *Regarding isomorphism classes of semisimple and simply connected affine group schemes over \mathbb{Z} :*

- (1) *there are two of type E_6 , namely $\mathbf{Isom}(\mathrm{Her}_3(C))$ and*
- (2) *there are two of type E_7 , namely $\mathbf{Aut}(Q(\mathrm{Her}_3(C)))$*

for $C = \mathrm{Zor}(\mathbb{Z})$ or \mathcal{O} .

Proof. Put \mathbf{G} for the simply connected Chevalley group scheme over \mathbb{Z} of type E_n , for $n = 6$ or 7 . By [Con, Remark 4.8], \mathbb{Z} forms of absolutely simple and simply connected \mathbb{Q} -group schemes are purely inner forms, i.e., are obtained by twisting \mathbf{G} by a class $\xi \in H^1(\mathbb{Z}, \mathbf{G})$.

For the split Albert algebra $J = \mathrm{Her}_3(\mathrm{Zor}(\mathbb{Z}))$, the natural inclusions $\mathbf{Aut}(J) \subset \mathbf{Isom}(J) \subset \mathbf{Aut}(Q(J))$ give maps $H^1(R, \mathbf{Aut}(J)) \rightarrow H^1(R, \mathbf{G})$ for every ring R , where the domain is in bijection with the isomorphism classes of Albert R -algebras. The groups in the statement we are aiming to prove are the image of $\mathrm{Her}_3(C)$ for $C = \mathrm{Zor}(\mathbb{Z})$ or \mathcal{O} . The two choices for C , i.e., the two groups in the statement, give non-isomorphic groups over \mathbb{R} , so it suffices to show that there are no others defined over \mathbb{Z} .

Now the compact real form of type E_n is not a pure inner form, so $G_\xi \times \mathbb{R}$ is not compact for all $\xi \in H^1(\mathbb{Z}, \mathbf{G})$. Therefore, the natural map $H^1(\mathbb{Z}, \mathbf{G}) \rightarrow H^1(\mathbb{Q}, \mathbf{G})$ is a bijection by [Har67, Satz 4.2.4]. The natural map $H^1(\mathbb{Q}, \mathbf{G}) \rightarrow H^1(\mathbb{R}, \mathbf{G})$ is also a bijection, a fact we have already used in the proof of Proposition 13.2. Since $H^1(\mathbb{R}, \mathbf{G})$ has two elements — see [BoroE], [BoroT, esp. §15], or [AdT, Table 3] — we have produced both elements of $H^1(\mathbb{Z}, \mathbf{G})$. \square

The proof provides the following corollary.

Corollary 17.2. *There are two isomorphism classes of FT systems over \mathbb{Z} , namely $Q(\text{Her}_3(C))$ for $C = \text{Zor}(\mathbb{Z})$ or \mathcal{O} .* \square

We have addressed now all the simple types that are usually called “exceptional”, apart from E_8 . A classification of \mathbb{Z} -groups of type E_8 like Proposition 17.1 appears currently out of reach, because among those group schemes \mathbf{G} over \mathbb{Z} such that $\mathbf{G} \times \mathbb{R}$ is the compact group of type E_8 , there are at least 13,935 distinct isomorphism classes [Gro96, Prop. 5.3]. Among those \mathbf{G} over \mathbb{Z} of type E_8 such that $\mathbf{G} \times \mathbb{R}$ is not compact, the same argument as in the proof of Proposition 17.1 shows that there are two isomorphism classes.

REFERENCES

- [ABB14] A. Auel, M. Bernardara, and M. Bolognesi, *Fibrations in complete intersections of quadrics, clifford algebras, derived categories, and rationality problems*, J. Math. Pures Appl. **102** (2014), 249–291. [5](#)
- [AG19] S. Alsaody and P. Gille, *Isotopes of octonion algebras, \mathbf{G}_2 -torsors and triality*, Adv. Math. **343** (2019), 864–909. [16](#), [20](#)
- [AHW19] A. Asok, M. Hoyois, and M. Wendt, *Generically split octonion algebras and \mathbb{A}^1 -homotopy theory*, Algebra & Number Theory **13** (2019), no. 3, 695–747. [20](#)
- [AJ57] A.A. Albert and N. Jacobson, *On reduced exceptional simple Jordan algebras*, Ann. of Math. (2) **66** (1957), 400–417, (= Jacobson, Oe. 58). [19](#)
- [Alb42] A.A. Albert, *Non-associative algebras. I. Fundamental concepts and isotopy*, Annals Math. (2) **43** (1942), 687–707. [20](#)
- [Alb58] ———, *A construction of exceptional Jordan division algebras*, Annals of Mathematics **67** (1958), 1–28. [24](#)
- [Als21] S. Alsaody, *Albert algebras over rings and related torsors*, Canad. J. Math. **73** (2021), no. 3, 875–898. [2](#), [14](#), [21](#), [25](#)
- [Asc88] M. Aschbacher, *Some multilinear forms with large isometry groups*, Geometriae Dedicata **25** (1988), 417–465. [31](#)
- [AT18] J. Adams and O. Taïbi, *Galois and Cartan cohomology of real groups*, Duke Math. J. **167** (2018), no. 6, 1057–1097. [19](#), [32](#)
- [BC12] P. Balmer and B. Calmès, *Bases of total Witt groups and lax-similitude*, J. Algebra Appl. **11** (2012), no. 3, 1250045. [5](#)
- [BDF⁺14] L. Borsten, M.J. Duff, S. Ferrara, A. Marrani, and W. Rubens, *Explicit orbit classification of reducible Jordan algebras and Freudenthal triple systems*, Comm. Math. Phys. **325** (2014), 17–39. [31](#)
- [BE16] M. Borovoi and Z. Evenor, *Real homogeneous spaces, Galois cohomology, and Reeder puzzles*, J. Algebra **467** (2016), 307–365. [19](#), [32](#)
- [BK66] H. Braun and M. Koecher, *Jordan-Algebren*, Springer-Verlag, Berlin, 1966. [23](#)
- [Bou88] N. Bourbaki, *Algebra II: Chapters 4–7*, Springer-Verlag, 1988. [5](#), [12](#), [28](#)
- [Bro69] R.B. Brown, *Groups of type E_7* , J. Reine Angew. Math. **236** (1969), 79–102. [27](#), [28](#), [30](#)
- [BT21] M. Borovoi and D.A. Timashev, *Galois cohomology of real semisimple groups via Kac labelings*, Transf. Groups **26** (2021), no. 2, 443–477. [32](#)
- [CF15] B. Calmès and J. Fasel, *Groupes classiques*, Panoramas & Synthèses **46** (2015), 1–133. [3](#), [5](#), [25](#)
- [CG06] M. Carr and S. Garibaldi, *Geometries, the principle of duality, and algebraic groups*, Expo. Math. **24** (2006), 195–234. [15](#)
- [CG21] M. Chayet and S. Garibaldi, *A class of continuous non-associative algebras arising from algebraic groups including E_8* , Forum of Mathematics: Sigma **9** (2021), no. e6. [2](#)

- [Cle03] J.-L. Clerc, *Special prehomogeneous vector spaces associated to F_4, E_6, E_7, E_8 and simple Jordan algebras of rank 3*, J. Algebra **264** (2003), 98–128. [31](#)
- [Con14] B. Conrad, *Non-split reductive groups over \mathbb{Z}* , *Autour des Schémas en Groupes II*, Panoramas et synthèses, vol. 46, Société Mathématique de France, 2014, pp. 193–253. [2](#), [8](#), [16](#), [19](#), [24](#), [25](#), [31](#)
- [Cox46] H.S.M. Coxeter, *Integral Cayley numbers*, Duke Math. J. **13** (1946), 561–578. [8](#)
- [CS03] J.H. Conway and D.A. Smith, *On quaternions and octonions: their geometry, arithmetic, and symmetry*, A K Peters, Ltd., Natick, MA, 2003. [8](#)
- [DG70] M. Demazure and A. Grothendieck, *Schémas en groupes III: Structure des schémas en groupes reductifs*, Lecture Notes in Mathematics, vol. 153, Springer, 1970. [15](#), [17](#), [27](#)
- [DG02] P. Deligne and B.H. Gross, *On the exceptional series, and its descendants*, C. R. Math. Acad. Sci. Paris **335** (2002), no. 11, 877–881. [31](#)
- [Dic23] L.E. Dickson, *A new simple theory of hypercomplex integers*, Journal de Mathématiques Pures et Appliquées **29** (1923), 281–326. [8](#)
- [EG96] N.D. Elkies and B.H. Gross, *The exceptional cone and the Leech lattice*, Internat. Math. Res. Notices (1996), no. 14, 665–698. [2](#), [8](#), [23](#)
- [EKM08] R.S. Elman, N. Karpenko, and A. Merkurjev, *The algebraic and geometric theory of quadratic forms*, Colloquium Publications, vol. 56, Amer. Math. Soc., 2008. [24](#)
- [Fer72] J.C. Ferrar, *Strictly regular elements in Freudenthal triple systems*, Trans. Amer. Math. Soc. **174** (1972), 313–331. [29](#)
- [Fre54] H. Freudenthal, *Beziehungen der E_7 und E_8 zur Oktavenebene. I*, Nederl. Akad. Wetensch. Proc. Ser. A **57** (1954), 218–230. [30](#)
- [Fre85] ———, *Oktaven, Ausnahmegruppen und Oktavengeometrie*, Geom. Dedicata **19** (1985), 7–63, (Reprint of Utrecht Lecture Notes, 1951). [16](#)
- [Gar01] S. Garibaldi, *Structurable algebras and groups of type E_6 and E_7* , J. Algebra **236** (2001), no. 2, 651–691. [30](#), [31](#)
- [Gar09] ———, *Cohomological invariants: exceptional groups and spin groups*, Memoirs Amer. Math. Soc., no. 937, Amer. Math. Soc., Providence, RI, 2009, with an appendix by Detlev W. Hoffmann. [24](#), [26](#), [31](#)
- [GG21] B.H. Gross and S. Garibaldi, *Minuscule embeddings*, Indag. Math. **32** (2021), no. 5, 987–1004. [31](#)
- [Gil14] P. Gille, *Octonion algebras over rings are not determined by their norms*, Canad. Math. Bull. **57** (2014), no. 2, 303–309. [20](#)
- [Gil19] ———, *Groupes algébriques semi-simples en dimension cohomologique ≤ 2* , Lecture Notes in Math., no. 2238, Springer, 2019. [19](#)
- [Gir71] J. Giraud, *Cohomologie non abélienne*, Die Grundlehren der mathematischen Wissenschaften, vol. 179, Springer-Verlag, Berlin, 1971. [3](#)
- [Gro96] B.H. Gross, *Groups over \mathbb{Z}* , Invent. Math. **124** (1996), no. 1-3, 263–279. [2](#), [24](#), [32](#)
- [Har66] G. Harder, *Über die Galoiskohomologie halbeinfacher Matrizenruppen. II*, Math. Z. **92** (1966), 396–415. [23](#)
- [Har67] ———, *Halbeinfache Gruppenschemata über Dedekindringen*, Invent. Math. **4** (1967), 165–191. [19](#), [20](#), [32](#)
- [Hel12] F. Helenius, *Freudenthal triple systems by root system methods*, J. Algebra **357** (2012), 116–137. [31](#)
- [Hij63] H. Hijikata, *A remark on the groups of type G_2 and F_4* , J. Math. Soc. Japan **15** (1963), 159–164. [17](#)
- [Jac61] N. Jacobson, *Some groups of transformations defined by Jordan algebras. III*, J. Reine Angew. Math. **207** (1961), 61–85, (= Coll. Math. Papers 65). [21](#)
- [Jac68] ———, *Structure and representations of Jordan algebras*, Coll. Pub., vol. 39, Amer. Math. Soc., Providence, RI, 1968. [2](#), [9](#), [24](#)
- [Jac69] ———, *Lectures on quadratic Jordan algebras*, Tata Institute of Fundamental Research, Bombay, 1969, Tata Institute of Fundamental Research Lectures on Mathematics, No. 45. [10](#)
- [Jac71] ———, *Exceptional Lie algebras*, Lecture notes in pure and applied mathematics, vol. 1, Marcel-Dekker, New York, 1971. [16](#), [21](#), [25](#)
- [KMRT98] M.-A. Knus, A.S. Merkurjev, M. Rost, and J.-P. Tignol, *The book of involutions*, Colloquium Publications, vol. 44, Amer. Math. Soc., 1998. [2](#), [17](#)
- [Knu91] M.-A. Knus, *Quadratic and Hermitian forms over rings*, Springer-Verlag, Berlin, 1991. [5](#), [7](#), [8](#)
- [KO74] M.-A. Knus and M. Ojanguren, *Théorie de la descente et algèbres d’Azumaya*, Lecture Notes in Mathematics, vol. 389, Springer-Verlag, 1974. [5](#), [14](#)
- [Kru02] S. Krutevich, *On a canonical form of a 3×3 Hermitian matrix over the ring of integral split octonions*, J. Algebra **253** (2002), 276–295. [21](#)

- [Kru07] ———, *Jordan algebras, exceptional groups, and Bhargava composition*, J. Algebra **314** (2007), no. 2, 924–977. [27](#), [30](#), [31](#)
- [Loo06] O. Loos, *Generically algebraic Jordan algebras over commutative rings*, J. Algebra **297** (2006), 474–529. [3](#), [18](#)
- [LPR08] O. Loos, H.P. Petersson, and M.L. Racine, *Inner derivations of alternative algebras over commutative rings*, Algebra Number Theory **2** (2008), no. 8, 927–968. [2](#), [8](#)
- [Lur01] J. Lurie, *On simply laced Lie algebras and their minuscule representations*, Comment. Math. Helv. **76** (2001), 515–575. [27](#), [30](#)
- [Luz13] A.Yu. Luzgarev, *Fourth-degree invariants for $G(E_7, R)$ not depending on the characteristic*, Vestnik St. Petersburg University. Mathematics **46** (2013), no. 1, 29–34. [31](#)
- [Mah42] K. Mahler, *On ideals in the Cayley-Dickson algebra*, Proc. Roy. Irish Acad. Sect. A. **48** (1942), 123–133. [14](#)
- [McC66] K. McCrimmon, *A general theory of Jordan rings*, Proc. Nat. Acad. Sci. U.S.A. **56** (1966), 1072–1079. [8](#), [9](#), [15](#)
- [McC69] ———, *The Freudenthal-Springer-Tits constructions of exceptional Jordan algebras*, Trans. Amer. Math. Soc. **139** (1969), 495–510. [9](#), [10](#), [11](#), [20](#)
- [McC71a] ———, *Homotopes of alternative algebras*, Math. Ann. **191** (1971), 253–262. [13](#)
- [McC71b] ———, *Inner ideals in quadratic Jordan algebras*, Trans. Amer. Math. Soc. **159** (1971), 445–468. [15](#)
- [McC85] ———, *Nonassociative algebras with scalar involution*, Pacific J. Math. **116** (1985), no. 1, 85–109. [11](#)
- [McC04] ———, *A taste of Jordan algebras*, Universitext, Springer-Verlag, New York, 2004. [9](#)
- [Mey68] K. Meyberg, *Eine Theorie der Freudenthalschen Tripelsysteme. I, II*, Nederl. Akad. Wetensch. Proc. Ser. A 71 = Indag. Math. **30** (1968), 162–190. [28](#)
- [Mil17] J.S. Milne, *Algebraic groups: the theory of group schemes of finite type over a field*, Cambridge studies in Advanced Math., vol. 170, Cambridge University Press, 2017. [16](#)
- [MW19] B. Mühlherr and R.M. Weiss, *Freudenthal triple systems in arbitrary characteristic*, J. Algebra **520** (2019), 237–275. [31](#)
- [MZ88] K. McCrimmon and E. Zel’manov, *The structure of strongly prime quadratic Jordan algebras*, Adv. in Math. **69** (1988), no. 2, 133–222. [2](#)
- [Pet93] H.P. Petersson, *Composition algebras over algebraic curves of genus zero*, Trans. Amer. Math. Soc. **337** (1993), no. 1, 473–493. [6](#)
- [Pet19] ———, *A survey on Albert algebras*, Transf. Groups **24** (2019), no. 1, 219–278. [2](#), [13](#), [14](#), [18](#)
- [Pet21] ———, *Norm equivalences and ideals of composition algebras*, Münster J. of Math. **14** (2021), 283–293. [14](#)
- [PR85] H.P. Petersson and M.L. Racine, *Radicals of Jordan algebras of degree 3*, Radical theory (Eger, 1982), Colloq. Math. Soc. János Bolyai, vol. 38, North-Holland, Amsterdam, 1985, pp. 349–377. [11](#)
- [PR86] ———, *Jordan algebras of degree 3 and the Tits process*, J. Algebra **98** (1986), no. 1, 211–243. [9](#), [10](#), [24](#)
- [PR94] V.P. Platonov and A. Rapinchuk, *Algebraic groups and number theory*, Academic Press, Boston, 1994. [19](#), [23](#)
- [PR96] H.P. Petersson and M.L. Racine, *An elementary approach to the Serre-Rost invariant of Albert algebras*, Indag. Math. (N.S.) **7** (1996), no. 3, 343–365. [24](#)
- [Rac77] M.L. Racine, *Point spaces in exceptional quadratic Jordan algebras*, J. Algebra **46** (1977), 22–36. [15](#)
- [Rob63] N. Roby, *Lois polynomes et lois formelles in théorie des modules*, Ann. Sci. École Norm. Sup. (3) **80** (1963), no. 3, 213–348. [3](#), [5](#), [10](#)
- [Röh93] G. Röhrle, *On extraspecial parabolic subgroups*, Linear algebraic groups and their representations (Los Angeles, CA, 1992), Contemp. Math., vol. 153, Amer. Math. Soc., Providence, RI, 1993, pp. 143–155. [31](#)
- [Ros91] M. Rost, *A (mod 3) invariant for exceptional Jordan algebras*, C. R. Acad. Sci. Paris Sér. I Math. **313** (1991), 823–827. [24](#)
- [Sch94] R. Schafer, *An introduction to nonassociative algebras*, Dover, 1994. [6](#)
- [Sel63] G.B. Seligman, *On the split exceptional Lie algebra E_7* , dittoed notes, 28 pages, 1963. [30](#)
- [Ser02] J-P. Serre, *Galois cohomology*, Springer-Verlag, 2002, originally published as *Cohomologie galoisienne* (1965). [3](#), [16](#), [19](#)
- [Ses77] C.S. Seshadri, *Geometric reductivity over arbitrary base*, Adv. Math. **26** (1977), 225–274. [29](#)

- [Spr62] T.A. Springer, *Characterization of a class of cubic forms*, *Nederl. Akad. Wetensch.* **65** (1962), 259–265. [26](#)
- [Spr73] ———, *Jordan algebras and algebraic groups*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol. 75, Springer-Verlag, 1973. [16](#), [25](#)
- [Spr06] ———, *Some groups of type E_7* , *Nagoya Math. J.* **182** (2006), 259–284. [28](#), [31](#)
- [Sta18] The Stacks Project Authors, *Stacks project*, <https://stacks.math.columbia.edu>, 2018. [5](#), [12](#)
- [SV00] T.A. Springer and F.D. Veldkamp, *Octonions, Jordan algebras and exceptional groups*, Springer-Verlag, Berlin, 2000. [16](#), [18](#), [19](#), [25](#)
- [Tit90] J. Tits, *Strongly inner anisotropic forms of simple algebraic groups*, *J. Algebra* **131** (1990), 648–677. [31](#)
- [vdBS59] F. van der Blij and T.A. Springer, *The arithmetics of octaves and of the group G_2* , *Nederl. Akad. Wetensch.* **62** (1959), 406–418. [24](#)
- [Voi21] J. Voight, *Quaternion algebras*, Springer, 2021. [8](#)
- [Wat79] W.C. Waterhouse, *Introduction to affine group schemes*, *Graduate Texts in Mathematics*, vol. 66, Springer, 1979. [3](#)
- [Wei60] A. Weil, *Algebras with involution and the classical groups*, *J. Indian Math. Soc.* **24** (1960), 589–623. [2](#)